



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C12Q 1/68, C12P 19/34, C12N 1/21,</b> <b>9/16, 15/63, 15/70, C07H 21/02, 21/04</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/29482</b> <b>(43) International Publication Date:</b> 22 December 1994 (22.12.94)
<b>(21) International Application Number:</b> PCT/US94/06253 <b>(22) International Filing Date:</b> 6 June 1994 (06.06.94)  <b>(30) Priority Data:</b> 08/073,384                      4 June 1993 (04.06.93)                      US  <b>(71) Applicant:</b> THIRD WAVE TECHNOLOGIES, INC. [US/US]; 2800 Fish Hatchery Road, Madison, WI 53711-5368 (US).  <b>(72) Inventors:</b> DAHLBERG, James, E.; 1119 Merrill Springs Road, Madison, WI 53705-1316 (US). LYAMICHEV, Victor, I.; 2221 Post Road #5, Madison, WI 53713 (US). BROW, Mary, Ann, D.; 5905 Hammersley Road, Madison, WI 53711 (US).  <b>(74) Agents:</b> CARROLL, Peter, G. et al.; Haverstock, Medlen & Carroll, Suite 2200, 220 Montgomery Street, San Francisco, CA 94104 (US).		<b>(81) Designated States:</b> AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> 5' NUCLEASES DERIVED FROM THERMOSTABLE DNA POLYMERASE		
<b>(57) Abstract</b>  A means for cleaving a nucleic acid cleavage structure in a site-specific manner is disclosed. A cleaving enzyme having 5' nuclease activity without interfering nucleic acid synthetic ability is employed as the basis of a novel method of detection of specific nucleic acid sequences.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## 5' NUCLEASES DERIVED FROM THERMOSTABLE DNA POLYMERASE

This is a Continuation-In-Part Application of Application Serial No. 08/073,384, filed June 4, 1993, which is a Continuation-In-Part Application of Application Serial No. 07/986,330, filed December 12, 1992.

### 5 FIELD OF THE INVENTION

The present invention relates to means for cleaving a nucleic acid cleavage structure in a site-specific manner. In particular, the present invention relates to a cleaving enzyme having 5' nuclease activity without interfering nucleic acid synthetic ability.

### 10 BACKGROUND OF THE INVENTION

The detection of specific nucleic acid sequences has been utilized to diagnose the presence of viral or bacterial nucleic acid sequences indicative of an infection, the presence of variants or alleles of mammalian genes associated with disease and the identification of the source of nucleic acids found in forensic  
15 samples and in paternity determinations.

The detection of specific nucleic acid sequences has been achieved typically by hybridization. Hybridization methods involve the annealing of a complementary sequence to the target nucleic acid (the sequence to be detected). The ability of two polymers of nucleic acid containing complementary sequences to find each  
20 other and anneal through base pairing interaction is a well-recognized phenomenon. The initial observations of the "hybridization" process by Marmur and Lane, *Proc. Natl. Acad. Sci. USA* 46:453 (1960) and Doty *et al.*, *Proc. Natl. Acad. Sci. USA* 46:461 (1960) have been followed by the refinement of this process into an essential tool of modern biology.

25 Initial hybridization studies, such as those performed by Hayashi *et al.*, *Proc. Natl. Acad. Sci. USA* 50:664 (1963), were formed in solution. Further development led to the immobilization of the target DNA or RNA on solid supports. With the discovery of specific restriction endonucleases by Smith and Wilcox, *J. Mol. Biol.* 51:379 (1970), it became possible to isolate discrete

fragments of DNA. Utilization of immobilization techniques, such as those described by Southern, *J. Mol. Biol.* 98:503 (1975), in combination with restriction enzymes, has allowed for the identification by hybridization of single copy genes among a mass of fractionated, genomic DNA.

5 In spite of the progress made in hybridization methodology, a number of problems have prevented the wide scale use of hybridization as a tool in human diagnostics. Among the more formidable problems are: 1) the inefficiency of hybridization; 2) the low concentration of specific target sequences in a mixture of genomic DNA; and 3) the hybridization of only partially complementary probes  
10 and targets.

### 1. Inefficient Hybridization

It is experimentally observed that only a fraction of the possible number of probe-target complexes are formed in a hybridization reaction. This is particularly true with short oligonucleotide probes (less than 100 bases in length). There are  
15 three fundamental causes: a) hybridization cannot occur because of secondary and tertiary structure interactions; b) strands of DNA containing the target sequence have rehybridized (reannealed) to their complementary strand; and c) some target molecules are prevented from hybridization when they are used in hybridization formats that immobilize the target nucleic acids to a solid surface.

20 Even where the sequence of a probe is completely complementary to the sequence of the target, *i.e.*, the target's primary structure, the target sequence must be made accessible to the probe via rearrangements of higher-order structure. These higher-order structural rearrangements may concern either the secondary structure or tertiary structure of the molecule. Secondary structure is determined  
25 by intramolecular bonding. In the case of DNA or RNA targets this consists of hybridization within a single, continuous strand of bases (as opposed to hybridization between two different strands). Depending on the extent and position of intramolecular bonding, the probe can be displaced from the target sequence preventing hybridization.

30 Solution hybridization of oligonucleotide probes to denatured double-stranded DNA is further complicated by the fact that the longer complementary



target strands can renature or reanneal. Again, hybridized probe is displaced by this process. This results in a low yield of hybridization (low "coverage") relative to the starting concentrations of probe and target.

5 The immobilization of target nucleic acids to solid surfaces such as nylon or nitrocellulose is a common practice in molecular biology. Immobilization formats eliminate the reassociation problem that can occur between complementary strands of target molecules, but not the problems associated with secondary structure effects. However, these mixed phase formats (*i.e.*, Southern hybridization or dot blot hybridization) require time consuming fixation procedures. The hybridization  
10 reaction itself is kinetically much slower than a solution phase hybridization reaction. Together, the fixation and hybridization procedures require a minimum of several hours to several days to perform. Additionally, the standard immobilization procedures are often inefficient and result in the attachment of many of the target molecules to multiple portions on the solid surface, rendering them incapable of  
15 subsequent hybridization to probe molecules. Overall, these combined effects result in just a few percent of the initial target molecules being bound by probes in a hybridization reaction.

## 2. Low Target Sequence Concentration

In laboratory experiments, purified probes and targets are used. The  
20 concentrations of these probes and targets, moreover, can be adjusted according to the sensitivity required. By contrast, the goal in the application of hybridization to medical diagnostics is the detection of a target sequence from a mixture of genomic DNA. Usually the DNA fragment containing the target sequence is in relatively low abundance in genomic DNA. This presents great technical difficulties; most  
25 conventional methods that use oligonucleotide probes lack the sensitivity necessary to detect hybridization at such low levels.

One attempt at a solution to the target sequence concentration problem is the amplification of the detection signal. Most often this entails placing one or more labels on an oligonucleotide probe. In the case of non-radioactive labels,  
30 even the highest affinity reagents have been found to be unsuitable for the detection of single copy genes in genomic DNA with oligonucleotide probes. See Wallace

*et al.*, *Biochimie* 67:755 (1985). In the case of radioactive oligonucleotide probes, only extremely high specific activities are found to show satisfactory results. See Studencki and Wallace, *DNA* 3:1 (1984) and Studencki *et al.*, *Human Genetics* 37:42 (1985).

5 Polymerase chain reaction (PCR) technology provides an alternate approach to the problems of low target sequence concentration. PCR can be used to directly increase the concentration of the target prior to hybridization. In U.S. Patents Nos. 4,683,195 and 4,683,202, Mullis *et al.* describe a method for increasing the concentration of a segment of target sequence in a mixture of genomic DNA  
10 without cloning or purification.

This process for amplifying the target sequence consists of introducing a molar excess of two oligonucleotide primers to the DNA mixture containing the desired target sequence. The two primers are complementary to their respective strands of the double-stranded sequence. The mixture is denatured and then  
15 allowed to hybridize. Following hybridization, the primers are extended with polymerase so as to form complementary strands. The steps of denaturation, hybridization, and polymerase extension can be repeated as often as needed to obtain relatively high concentration of a segment of the desired target sequence. The length of the segment of the desired target sequence is determined by the  
20 relative positions of the primers with respect to each other, and, therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to by the inventors as the "Polymerase Chain Reaction" (or PCR). Because the desired segment of the target sequence become the dominant sequences (in terms of concentration) in the mixture, they are said to be "PCR-  
25 amplified."

However the PCR process is susceptible to the production of non-target fragments during the amplification process. Spurious extension of primers at partially complementary regions occurs during PCR reactions. Factors influencing the specificity of the amplification process include: a) the concentration of the  
30 target sequence in the DNA to be analyzed; b) the concentration of the  $Mg^{++}$ , polymerase enzyme and primers; c) the number of cycles of amplification performed; and d) the temperatures and times used at the various steps in the

amplification process [*PCR Technology - Principles and Applications for DNA Amplification* (H.A. Erlich, Ed.), Stockton Press, New York, pp. 7-16 (1989)].

When the specific target sequence is present in low concentration in the sample DNA more non-target fragments are produced. Low target concentration is often the norm in clinical samples where the target may be present as a single copy in the genome or where very little viral DNA is present as in HIV infections.

Because amplification products are produced which do not represent the specific target sequence to be detected, the products of a PCR reaction must be analyzed using a probe specific for the target DNA. The detection of specific amplification products has been accomplished by the hybridization of a probe specific for the target sequence to the reaction products immobilized upon a solid support. Such a detection method is cumbersome and is subject to the same problems associated with the detection of any target molecule by hybridization as discussed above.

A non-hybridization based detection assay for specific PCR products has been described by Holland *et al.*, *Proc. Natl. Acad. Sci. USA* 88:7276 (1991). In this detection system, the 5' nuclease activity of wild type DNA polymerase from *Thermus aquaticus* ("DNAP<sub>Taq</sub>") is used to generate a specific detectable product concomitantly with amplification. An oligonucleotide probe specific for the target DNA is labeled on the 5' end and added to the PCR reaction along with the unlabelled primers used for extension of the target to be amplified. The 5' nuclease activity of the DNAP<sub>Taq</sub> cleaves the labeled probe annealed to the target DNA before the extension of the primer is complete, generating a smaller fragment of the probe. This detection system requires that amplification be performed upon the sample to produce the specific detection product. This is slow and requires cumbersome equipment.

A minimum of 100 starting copies (*i.e.*, copy number prior to amplification) of target DNA were used in this detection system; it is not clear whether fewer starting copies of target DNA will yield detectable results using this method. Very low copy number may be a problem for some clinical samples where very little DNA is obtained due to restrictions on sample size (blood from neonates or fetuses, forensic samples, *etc.*).

While such an assay is an improvement over earlier hybridization detection methods, it still requires that a PCR reaction be performed upon the sample and it possesses certain inherent problems. One such problem is that this system requires that the detection probe must bind to the target DNA before primer extension occurs. If extension occurs first, the probe binding site will be unavailable and no digestion of the probe will occur and therefore no detectable signal will be produced. To overcome this problem the user must vary the relative amounts of primer and probe or manipulate the sequence and length of the probe. The need for such optimization may prove too burdensome for clinical laboratories.

### 3. Partial Complementarity

Hybridization, regardless of the method used, requires some degree of complementarity between the sequence being assayed (the target sequence) and the fragment of DNA used to perform the test (the probe). (Of course, one can obtain binding without any complementarity but this binding is nonspecific and to be avoided.) For many diagnostic applications, it is not important to determine whether the hybridization represents complete or partial complementarity. For example, where it is desired to detect simply the presence or absence of pathogen DNA (such as from a virus, bacterium, fungi, mycoplasma, protozoan) it is only important that the hybridization method ensures hybridization when the relevant sequence is present; conditions can be selected where both partially complementary probes and completely complementary probes will hybridize. Other diagnostic applications, however, may require that the method of hybridization distinguish between variant target sequences. For example, it may be of interest that a particular allelic variant of a pathogen is present. These normal and variant sequences may differ in one or more bases.

There are other applications that may require that the hybridization method distinguish between partial and complete complementarity. It may be of interest to detect genetic polymorphisms. Human hemoglobin is composed, in part, of four polypeptide chains. Two of these chains are identical chains of 141 amino acids (alpha chains) and two of these chains are identical chains of 146 amino acids (beta chains). The gene encoding the beta chain is known to exhibit polymorphism. The

normal allele encodes a beta chain having glutamic acid at the sixth position. The mutant allele encodes a beta chain having valine at the sixth position. This difference in amino acids has a profound (most profound when the individual is homozygous for the mutant allele) physiological impact known clinically as sickle cell anemia. It is well known that the genetic basis of the amino acid change involves a single base difference between the normal allele DNA sequence and the mutant allele DNA sequence.

Unless combined with other techniques (such as restriction enzyme analysis), hybridization methods that allow for the same level of hybridization in the case of both partial as well as complete complementarity are unsuited for such applications; the probe will hybridize to both the normal and variant target sequence.

Methods have been devised to enable discrimination between partial and complete complementarity. One approach is to take advantage of the temperature requirements of the specific hybridization under study. In typical melting curve experiments, such as those described by Wallace *et al.*, *Nucl. Acids Res.* 6:3543 (1979) and *Nucl. Acids Res.* 9:879 (1981), an immobilized probe-target complex is washed at increasing temperatures under non-equilibrium conditions. It is observed that partially complementary probe-target complexes display a lower thermal stability as compared to completely complementary probe-target complexes. This difference can be used, therefore, to determine whether the probe has hybridized to the partially complementary or the completely complementary target sequence.

Conventional methods that utilize the temperature dependant nature of hybridization are artful. The application of this method for the discrimination of single base mutations in human genomic targets is limited to the use of short oligonucleotide probes where the hybridization interaction with the target sequence is in the size range of 17 bases to 25 bases in length. The lower length limit is determined by the random probability of having a complement to the probe in the human genome, which is greater than 1 for a random 16 base pair interaction, but less than 1 for interactions 17 bases or longer in length. The upper limit is one of practicality. It is difficult to differentiate single base mismatches on the basis of thermal stability for interactions longer than 25 bases in length. These



conventional methods are, unfortunately also time consuming. Probe concentrations in these experiments are approximately  $1-5 \times 10^{-10}$ M. These concentrations are empirically derived; they minimize the use of probe and simultaneously provide sufficient discrimination to distinguish single copy genes utilizing probes of approximately 20 nucleotides in length. Hybridization times are two to ten hours at these concentrations. After hybridization, several washes of varying stringency are employed to remove excess probe, non-specifically bound probe, and probe bound to partially complementary sequences in the target genome. Careful control of these wash steps is necessary, since the signal (specifically bound probe) to noise (non-specifically bound probe) ratio of the experiment is ultimately determined by the wash procedures.

No detection method heretofore described has solved all three of the problems discussed above. The PCR process solves the problem of low target concentration. However, the specific detection of PCR products by any hybridization method is subject to the same problems associated with the detection of any target molecules. The detection of single base differences between PCR targets was initially accomplished through the use of a restriction enzyme analysis of the hybridization complexes formed between oligonucleotide probes and PCR targets. This technique is limited by that fact that restriction enzymes do not exist for all sequences. More recent studies have achieved discrimination without restriction enzymes, however these studies have involved the inefficient immobilization of target nucleic acids to solid surfaces [dot blot hybridization; Saiki *et al.*, *Nature* 324:163 (1986)].

Another method for the detection of allele-specific variants is disclosed by Kwok *et al.*, *Nucl. Acids Res.* 18:999 (1990). This method is based upon the fact that it is difficult for a DNAP to synthesize a DNA strand when there is a mismatch between the template strand and the primer. The mismatch acts to prevent the extension thereby preventing the amplification of a target DNA that is not perfectly complementary to the primer used in a PCR reaction. While an allele-specific variant may be detected by the use of a primer that is perfectly matched with only one of the possible alleles, this method of detection is artful and has limitations. Particularly troublesome is the fact that the base composition of

the mismatch influences the ability to prevent extension across the mismatch. Certain mismatches do not prevent extension or have only a minimal effect.

An ideal method of detecting specific target DNAs would allow detection without the need to amplify the sample DNA first and would allow the detection of target sequences which are present in low copy numbers in the DNA sample. This ideal method would also allow the discrimination between variants of the target sequence such that single base variations between alleles of mammalian genes can be discerned.

One object of the present invention is to provide a method of detection of specific nucleic acid sequences that solves the above-named problems.

### SUMMARY OF THE INVENTION

The present invention relates to means for cleaving a nucleic acid cleavage structure in a site-specific manner. In one embodiment, the means for cleaving is a cleaving enzyme comprising 5' nucleases derived from thermostable DNA polymerases. These polymerases form the basis of a novel method of detection of specific nucleic acid sequences. The present invention contemplates use of the novel detection method for, among other uses, clinical diagnostic purposes.

In one embodiment, the present invention contemplates a DNA sequence encoding a DNA polymerase altered in sequence (*i.e.*, a "mutant" DNA polymerase) relative to the native sequence such that it exhibits altered DNA synthetic activity from that of the native (*i.e.*, "wild type") DNA polymerase. It is preferred that the encoded DNA polymerase is altered such that it exhibits reduced synthetic activity from that of the native DNA polymerase. In this manner, the enzymes of the invention are predominantly 5' nucleases and are capable of cleaving nucleic acids in a structure-specific manner in the absence of interfering synthetic activity.

Importantly, the 5' nucleases of the present invention are capable of cleaving linear duplex structures to create single discrete cleavage products. These linear structures are either 1) not cleaved by the wild type enzymes (to any significant degree), or 2) are cleaved by the wild type enzymes so as to create multiple products. This characteristic of the 5' nucleases has been found to be

consistent of enzymes derived in this manner from thermostable polymerases across eubacterial thermophilic species.

It is not intended that the invention be limited by the nature of the alteration necessary to render the polymerase synthesis deficient nor the extent of the deficiency. The present invention contemplates altered structure (primary,  
5 secondary, *etc.*) as well as native structure inhibited by synthesis inhibitors.

Where the structure is altered, it is not intended that the invention be limited by the means by which the structure of the polymerase is altered. In one embodiment, the alteration of the native DNA sequence comprises a change in a  
10 single nucleotide. In another embodiment, the alteration of the native DNA sequence comprises a deletion of one or more nucleotides. In yet another embodiment, the alteration of the native DNA sequence comprises an insertion of one or more nucleotides. In either of these cases, the change in DNA sequence may manifest itself in a change in amino acid sequence.

15 The present invention contemplates 5' nucleases from a variety of sources. The preferred 5' nucleases are thermostable. Thermostable 5' nucleases are contemplated as particularly useful in that they operate at temperatures where nucleic acid hybridization is extremely specific, allowing for allele-specific detection (including single-base mismatches). In one embodiment, the thermostable  
20 5' nucleases are selected from the group consisting of altered polymerases derived from the native polymerases of *Thermus aquaticus*, *Thermus flavus* and *Thermus thermophilus*.

As noted above, the present invention contemplates the use of altered polymerases in a detection method. In one embodiment, the present invention  
25 contemplates a method of detecting the presence of a specific target nucleic acid molecule comprising: a) providing: i) a cleavage means, ii) a target nucleic acid, iii) a first oligonucleotide complementary to a first portion of said target nucleic acid, iv) a first solid support having a second oligonucleotide, a region of which is complementary to a second portion of said target nucleic acid, said non-  
30 complementary region of said second oligonucleotide providing a single-stranded arm at its 5' end, a portion of said 5' arm comprising a first signal oligonucleotide, v) a plurality of "uncleaved" second solid supports each having a third



oligonucleotide, a region of which is complementary to said first signal oligonucleotide, the non-complementary region of said third oligonucleotide providing a single-stranded arm at its 5' end, a portion of said 5' arm comprising a second signal oligonucleotide, and vi) a plurality of "uncleaved" third solid supports each having a fourth oligonucleotide, a region of which is complementary to said second signal oligonucleotide, the non-complementary region of said fourth oligonucleotide providing a single-stranded arm at its 5' end, a portion of said 5' arm comprising said first signal oligonucleotide; b) mixing said cleavage means, said target nucleic acid, said first oligonucleotide and said second oligonucleotide under conditions wherein said first oligonucleotide and the 3' end of said second oligonucleotide are annealed to said target DNA sequence so as to create a first cleavage structure and cleavage of said first cleavage structure results in the liberating of said first signal oligonucleotide; d) reacting said liberated first signal oligonucleotide with one of said plurality of second solid supports under conditions such that said first signal oligonucleotide hybridizes to said complementary region of said third oligonucleotide to create a second cleavage structure and cleavage of said second cleavage structure results in the liberating of said second signal oligonucleotide and a "cleaved" second solid support; e) reacting said liberated second signal oligonucleotide with one of said plurality of third solid supports under conditions such that said second signal oligonucleotide hybridizes to said complementary region of said fourth oligonucleotide to create a third cleavage structure and cleavage of said third cleavage structure results in the liberating of a second molecule of said first signal oligonucleotide and a "cleaved" third solid support; and h) detecting the presence of said first and second signal oligonucleotides.

It is preferred that, after the hybridization of said first signal oligonucleotide and liberation of said second signal oligonucleotide, said first signal oligonucleotide is itself released from said "cleaved" second solid support and reacted with one of said plurality of "uncleaved" second solid supports. Similarly, it is preferred that, after the hybridization of said second signal oligonucleotide and liberation of said second molecule of said first signal oligonucleotide, said second signal oligonucleotide is itself released from said "cleaved" third solid support and reacted

with on of said plurality of "uncleaved" third solid supports. By the term "cleaved" and "uncleaved" it is not meant to indicate that the solid support (*e.g.*, a bead) is physically cleaved or uncleaved. Rather, it is meant to indicate the status of the oligonucleotide attached to the solid support.

5 By reference to a "solid support" it is not intended that the invention be limited to separate and discrete supports. For example, the invention contemplates a design where the oligos are on the same solid support, albeit separate in different regions. In one embodiment, the solid support is a microtiter well wherein the oligos are attached (*e.g.*, covalently) or coated (*e.g.*, non-covalently) in different  
10 regions of the well.

In a second embodiment, the present invention contemplates a method of detecting the presence of a specific target nucleic acid molecule comprising:  
a) providing: i) a target nucleic acid, ii) a first oligonucleotide complementary to a first portion of said target nucleic acid, and iii) a second oligonucleotide, a region  
15 of which is complementary to a second portion of said target nucleic acid, said non-complementary region of said second oligonucleotide providing a single-stranded arm at its 5' end; b) mixing said target nucleic acid, said first oligonucleotide and said second oligonucleotide under conditions wherein said first oligonucleotide and the 3' end of said second oligonucleotide are annealed to said  
20 target DNA sequence so as to create a first cleavage structure; c) providing a cleavage means under conditions such that cleavage of said first cleavage structure occurs preferentially at a site located within said second oligonucleotide in a manner dependent upon the annealing of said first and second oligonucleotides on said target nucleic acid, thereby liberating the single-stranded arm of said second  
25 oligonucleotide generating a third oligonucleotide; d) providing a first hairpin structure having a single-stranded 3' arm and a single-stranded 5' arm under conditions wherein said third oligonucleotide anneals to said single-stranded 3' arm of said first hairpin thereby creating a second cleavage structure; e) providing conditions under which cleavage of said second cleavage structure occurs by said  
30 cleavage means liberating the single-stranded 5' arm of said second cleavage structure so as to create reaction products comprising a fourth oligonucleotide and a first cleaved hairpin detection molecule; f) providing a second hairpin structure

having a single-stranded 3' arm and a single-stranded 5' arm under conditions wherein said fourth oligonucleotide anneals to the single-stranded 3' arm of said second hairpin thereby creating a third cleavage structure; g) providing conditions under which cleavage of said third cleavage structure occurs by said cleavage means, liberating the single-stranded 5' arm of said third cleavage structure so as to create reaction products comprising generating a fifth oligonucleotide identical in sequence to said third oligonucleotide and a second cleaved hairpin detection molecule; and h) detecting the presence of said first and second cleaved hairpin detection molecules.

In one embodiment, the detection method of the present invention allows the detection of specific target nucleic acid sequences present in a sample without the need to amplify the number of target copies prior to detection. In this embodiment, steps d) through g) of the method are repeated at least once.

In a preferred embodiment, the cleavage means comprises a cleavage enzyme comprising an altered thermostable DNA polymerase having reduced synthesis capability, *i.e.*, a 5' nuclease derived from a thermostable DNA polymerase. While a complete absence of synthesis is not required, it is desired that cleavage reactions occur in the absence of polymerase activity at a level where it interferes with the discrimination needed for detection.

While the cleavage of the second embodiment of the detection method of the present invention can be independent of the annealing of the oligonucleotides, it is preferred that the cleavage is primer-dependent. In other words, it is desired that the cleavage reactions of steps c), e) and g) will not occur absent the annealing of said first oligonucleotide, said third oligonucleotide and said fourth oligonucleotide, respectively.

While cleavage is site-specific, the present invention allows for cleavage at a variety of sites. In one embodiment, the cleavage reaction of step c) occurs within the annealed portion of said second oligonucleotide. In another embodiment, the cleavage reaction of step c) occurs within the non-annealed portion of said second oligonucleotide.

## DESCRIPTION OF THE DRAWINGS

Figure 1A provides a schematic of one embodiment of the detection method of the present invention.

5 Figure 1B provides a schematic of a second embodiment of the detection method of the present invention.

Figure 2 is a comparison of the nucleotide structure of the DNAP genes isolated from *Thermus aquaticus*, *Thermus flavus* and *Thermus thermophilus*; the consensus sequence is shown at the top of each row.

10 Figure 3 is a comparison of the amino acid sequence of the DNAP isolated from *Thermus aquaticus*, *Thermus flavus*, and *Thermus thermophilus*; the consensus sequence is shown at the top of each row.

Figures 4A-G are a set of diagrams of wild-type and synthesis-deficient DNAP $Taq$  genes.

Figure 5A depicts the wild-type *Thermus flavus* polymerase gene.

15 Figure 5B depicts a synthesis-deficient *Thermus flavus* polymerase gene.

Figure 6 depicts a structure which cannot be amplified using DNAP $Taq$ .

Figure 7 is a ethidium bromide-stained gel demonstrating attempts to amplify a bifurcated duplex using either DNAP $Taq$  or DNAP $Stf$ .

20 Figure 8 is an autoradiogram of a gel analyzing the cleavage of a bifurcated duplex by DNAP $Taq$  and lack of cleavage by DNAP $Stf$ .

Figures 9A-B are a set of autoradiograms of gels analyzing cleavage or lack of cleavage upon addition of different reaction components and change of incubation temperature during attempts to cleave a bifurcated duplex with DNAP $Taq$ .

25 Figures 10A-B are an autoradiogram displaying timed cleavage reactions, with and without primer.

Figures 11A-B are a set of autoradiograms of gels demonstrating attempts to cleave a bifurcated duplex (with and without primer) with various DNAPs.

30 Figures 12A shows the substrates and oligonucleotides used to test the specific cleavage of substrate DNAs targeted by pilot oligonucleotides.

Figure 12B shows an autoradiogram of a gel showing the results of cleavage reactions using the substrates and oligonucleotides shown Fig. 12A.

Figure 13A shows the substrate and oligonucleotide used to test the specific cleavage of a substrate RNA targeted by a pilot oligonucleotide.

Figure 13B shows an autoradiogram of a gel showing the results of a cleavage reaction using the substrate and oligonucleotide shown in Fig. 13A.

5        Figure 14 is a diagram of vector pTTQ18.

Figure 15 is a diagram of vector pET-3c.

Figure 16A-E depicts a set of molecules which are suitable substrates for cleavage by the 5' nuclease activity of DNAPs.

10        Figure 17 is an autoradiogram of a gel showing the results of a cleavage reaction run with synthesis-deficient DNAPs.

Figure 18 is an autoradiogram of a PEI chromatogram resolving the products of an assay for synthetic activity in synthesis-deficient DNAP*Taq* clones.

Figure 19A depicts the substrate molecule used to test the ability of synthesis-deficient DNAPs to cleave short hairpin structures.

15        Figure 19B shows an autoradiogram of a gel resolving the products of a cleavage reaction run using the substrate shown in Fig. 19A.

Figure 20A shows the A- and T-hairpin molecules used in the trigger/detection assay.

20        Figure 20B shows the sequence of the alpha primer used in the trigger/detection assay.

Figure 20C shows the structure of the cleaved A- and T-hairpin molecules.

Figure 20D depicts the complementarity between the A- and T-hairpin molecules.

25        Figure 21 provides the complete 206-mer duplex sequence employed as a substrate for the 5' nucleases of the present invention

Figures 22A and B show the cleavage of linear nucleic acid substrates (based on the 206-mer of Figure 21) by wild type DNAPs and 5' nucleases isolated from *Thermus aquaticus* and *Thermus flavus*.

30        Figure 23 provides a detailed schematic corresponding to the of one embodiment of the detection method of the present invention.

Figure 24 shows the propagation of cleavage of the linear duplex nucleic acid structures of Figure 23 by the 5' nucleases of the present invention.

Figure 25A shows the "nibbling" phenomenon detected with the DNAPs of the present invention.

Figure 25B shows that the "nibbling" of Figure 25A is 5' nucleolytic cleavage and not phosphatase cleavage.

5        Figure 26 demonstrates that the "nibbling" phenomenon is duplex dependent.

Figure 27 is a schematic showing how "nibbling" can be employed in a detection assay.

Figure 28 demonstrates that "nibbling" can be target directed.

## 10        **DESCRIPTION OF THE INVENTION**

The present invention relates to means for cleaving a nucleic acid cleavage structure in a site-specific manner. In particular, the present invention relates to a cleaving enzyme having 5' nuclease activity without interfering nucleic acid synthetic ability.

15        This invention provides 5' nucleases derived from thermostable DNA polymerases which exhibit altered DNA synthetic activity from that of native thermostable DNA polymerases. The 5' nuclease activity of the polymerase is retained while the synthetic activity is reduced or absent. Such 5' nucleases are capable of catalyzing the structure-specific cleavage of nucleic acids in the absence  
20        of interfering synthetic activity. The lack of synthetic activity during a cleavage reaction results in nucleic acid cleavage products of uniform size.

The novel properties of the polymerases of the invention form the basis of a method of detecting specific nucleic acid sequences. This method relies upon the amplification of the detection molecule rather than upon the amplification of the  
25        target sequence itself as do existing methods of detecting specific target sequences.

DNA polymerases (DNAPs), such as those isolated from *E. coli* or from thermophilic bacteria of the genus *Thermus*, are enzymes that synthesize new DNA strands. Several of the known DNAPs contain associated nuclease activities in addition to the synthetic activity of the enzyme.

30        Some DNAPs are known to remove nucleotides from the 5' and 3' ends of DNA chains [Kornberg, *DNA Replication*, W.H. Freeman and Co., San Francisco,



pp. 127-139 (1980)]. These nuclease activities are usually referred to as 5' exonuclease and 3' exonuclease activities, respectively. For example, the 5' exonuclease activity located in the N-terminal domain of several DNAPs participates in the removal of RNA primers during lagging strand synthesis during DNA replication and the removal of damaged nucleotides during repair. Some DNAPs, such as the *E. coli* DNA polymerase (DNAPEc1), also have a 3' exonuclease activity responsible for proof-reading during DNA synthesis (Kornberg, *supra*).

A DNAP isolated from *Thermus aquaticus*, termed *Taq* DNA polymerase (DNAPTaq), has a 5' exonuclease activity, but lacks a functional 3' exonucleolytic domain [Tindall and Kunkell, *Biochem.* 27:6008 (1988)]. Derivatives of DNAPEc1 and DNAPTaq, respectively called the Klenow and Stoffel fragments, lack 5' exonuclease domains as a result of enzymatic or genetic manipulations [Brutlag *et al.*, *Biochem. Biophys. Res. Commun.* 37:982 (1969); Erlich *et al.*, *Science* 252:1643 (1991); Setlow and Kornberg, *J. Biol. Chem.* 247:232 (1972)].

The 5' exonuclease activity of DNAPTaq was reported to require concurrent synthesis [Gelfand, *PCR Technology - Principles and Applications for DNA Amplification* (H.A. Erlich, Ed.), Stockton Press, New York, p. 19 (1989)]. Although mononucleotides predominate among the digestion products of the 5' exonucleases of DNAPTaq and DNAPEc1, short oligonucleotides ( $\leq 12$  nucleotides) can also be observed implying that these so-called 5' exonucleases can function endonucleolytically [Setlow, *supra*; Holland *et al.*, *Proc. Natl. Acad. Sci. USA* 88:7276 (1991)].

In WO 92/06200, Gelfand *et al.* show that the preferred substrate of the 5' exonuclease activity of the thermostable DNA polymerases is displaced single-stranded DNA. Hydrolysis of the phosphodiester bond occurs between the displaced single-stranded DNA and the double-helical DNA with the preferred exonuclease cleavage site being a phosphodiester bond in the double helical region. Thus, the 5' exonuclease activity usually associated with DNAPs is a structure-dependent single-stranded endonuclease and is more properly referred to as a 5' nuclease. Exonucleases are enzymes which cleave nucleotide molecules from the ends of the nucleic acid molecule. Endonucleases, on the other hand, are enzymes

which cleave the nucleic acid molecule at internal rather than terminal sites. The nuclease activity associated with some thermostable DNA polymerases cleaves endonucleolytically but this cleavage requires contact with the 5' end of the molecule being cleaved. Therefore, these nucleases are referred to as 5' nucleases.

5           When a 5' nuclease activity is associated with a eubacterial Type A DNA polymerase, it is found in the one-third N-terminal region of the protein as an independent functional domain. The C-terminal two-thirds of the molecule constitute the polymerization domain which is responsible for the synthesis of DNA. Some Type A DNA polymerases also have a 3' exonuclease activity  
10           associated with the two-third C-terminal region of the molecule.

          The 5' exonuclease activity and the polymerization activity of DNAPs have been separated by proteolytic cleavage or genetic manipulation of the polymerase molecule. To date thermostable DNAPs have been modified to remove or reduce the amount of 5' nuclease activity while leaving the polymerase activity intact.

15           The Klenow or large proteolytic cleavage fragment of DNAPEc1 contains the polymerase and 3' exonuclease activity but lacks the 5' nuclease activity. The Stoffel fragment of DNAP*Taq* lacks the 5' nuclease activity due to a genetic manipulation which deleted the N-terminal 289 amino acids of the polymerase molecule [Erlich *et al.*, *Science* 252:1643 (1991)]. WO 92/06200 describes a  
20           thermostable DNAP with an altered level of 5' to 3' exonuclease. U.S. Patent No. 5,108,892 describes a *Thermus aquaticus* DNAP without a 5' to 3' exonuclease. However, the art of molecular biology lacks a thermostable DNA polymerase with a lessened amount of synthetic activity.

          The present invention provides 5' nucleases derived from thermostable Type  
25           A DNA polymerases that retain 5' nuclease activity but have reduced or absent synthetic activity. The ability to uncouple the synthetic activity of the enzyme from the 5' nuclease activity proves that the 5' nuclease activity does not require concurrent DNA synthesis as was previously reported (Gelfand, *PCR Technology, supra*).

30           The description of the invention is divided into: I. Detection of Specific Nucleic Acid Sequences Using 5' Nucleases; II. Generation of 5' Nucleases Derived From Thermostable DNA Polymerases; III. Therapeutic Uses of 5'



Nucleases; and IV. Detection of Antigenic or Nucleic Acid Targets by a Dual Capture Assay. To facilitate understanding of the invention, a number of terms are defined below.

5 The term "gene" refers to a DNA sequence that comprises control and coding sequences necessary for the production of a polypeptide or precursor. The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired enzymatic activity is retained.

10 The term "wild-type" refers to a gene or gene product which has the characteristics of that gene or gene product when isolated from a naturally occurring source. In contrast, the term "modified" or mutant" refers to a gene or gene product which displays altered characteristics when compared to the wild-type gene or gene product. It is noted that naturally-occurring mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

15 The term "recombinant DNA vector" as used herein refers to DNA sequences containing a desired coding sequence and appropriate DNA sequences necessary for the expression of the operably linked coding sequence in a particular host organism. DNA sequences necessary for expression in procaryotes include a promoter, optionally an operator sequence, a ribosome binding site and possibly other sequences. Eucaryotic cells are known to utilize promoters, polyadenylation signals and enhancers.

20 The term "oligonucleotide" as used herein is defined as a molecule comprised of two or more deoxyribonucleotides or ribonucleotides, preferably more than three, and usually more than ten. The exact size will depend on many factors, which in turn depends on the ultimate function or use of the oligonucleotide. The oligonucleotide may be generated in any manner, including chemical synthesis, DNA replication, reverse transcription, or a combination thereof.

25 Because mononucleotides are reacted to make oligonucleotides in a manner such that the 5' phosphate of one mononucleotide pentose ring is attached to the 3' oxygen of its neighbor in one direction via a phosphodiester linkage, an end of an oligonucleotide is referred to as the "5' end" if its 5' phosphate is not linked to the 3' oxygen of a mononucleotide pentose ring and as the "3' end" if its 3' oxygen is

not linked to a 5' phosphate of a subsequent mononucleotide pentose ring. As used herein, a nucleic acid sequence, even if internal to a larger oligonucleotide, also may be said to have 5' and 3' ends.

5 When two different, non-overlapping oligonucleotides anneal to different regions of the same linear complementary nucleic acid sequence, and the 3' end of one oligonucleotide points towards the 5' end of the other, the former may be called the "upstream" oligonucleotide and the latter the "downstream" oligonucleotide.

10 The term "primer" refers to an oligonucleotide which is capable of acting as a point of initiation of synthesis when placed under conditions in which primer extension is initiated. An oligonucleotide "primer" may occur naturally, as in a purified restriction digest or may be produced synthetically.

15 A primer is selected to be "substantially" complementary to a strand of specific sequence of the template. A primer must be sufficiently complementary to hybridize with a template strand for primer elongation to occur. A primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment may be attached to the 5' end of the primer, with the remainder of the primer sequence being substantially complementary to the strand. Non-complementary bases or longer sequences can be interspersed into the  
20 primer, provided that the primer sequence has sufficient complementarity with the sequence of the template to hybridize and thereby form a template primer complex for synthesis of the extension product of the primer.

25 The complement of a nucleic acid sequence as used herein refers to an oligonucleotide which, when aligned with the nucleic acid sequence such that the 5' end of one sequence is paired with the 3' end of the other, is in "antiparallel association." Certain bases not commonly found in natural nucleic acids may be included in the nucleic acids of the present invention and include, for example, inosine and 7-deazaguanine. Complementarity need not be perfect; stable duplexes may contain mismatched base pairs or unmatched bases. Those skilled in the art of  
30 nucleic acid technology can determine duplex stability empirically considering a number of variables including, for example, the length of the oligonucleotide, base

composition and sequence of the oligonucleotide, ionic strength and incidence of mismatched base pairs.

5 Stability of a nucleic acid duplex is measured by the melting temperature, or " $T_m$ ." The  $T_m$  of a particular nucleic acid duplex under specified conditions is the temperature at which half of the base pairs have disassociated.

The term "probe" as used herein refers to a labeled oligonucleotide which forms a duplex structure with a sequence in another nucleic acid, due to complementarity of at least one sequence in the probe with a sequence in the other nucleic acid.

10 The term "label" as used herein refers to any atom or molecule which can be used to provide a detectable (preferably quantifiable) signal, and which can be attached to a nucleic acid or protein. Labels may provide signals detectable by fluorescence, radioactivity, colorimetry, gravimetry, X-ray diffraction or absorption, magnetism, enzymatic activity, and the like.

15 The term "cleavage structure" as used herein, refers to a nucleic acid structure which is a substrate for cleavage by the 5' nuclease activity of a DNAP.

The term "cleavage means" as used herein refers to any means which is capable of cleaving a cleavage structure in a specific manner. The cleavage means may include native DNAPs having 5' nuclease activity, and, more specifically,  
20 modified DNAPs having 5' nuclease but lacking synthetic activity.

The term "liberating" as used herein refers to the release of a nucleic acid fragment from a larger nucleic acid fragment, such as an oligonucleotide, by the action of a 5' nuclease such that the released fragment is no longer covalently attached to the remainder of the oligonucleotide.

25 The term "substrate strand" as used herein, means that strand of nucleic acid in a cleavage structure in which the cleavage mediated by the 5' nuclease activity occurs.

The term "template strand" as used herein, means that strand of nucleic acid in a cleavage structure which is at least partially complementary to the substrate strand and which anneals to the substrate strand to form the cleavage structure.  
30

The term " $K_m$ " as used herein refers to the Michaelis-Menten constant for an enzyme and is defined as the concentration of the specific substrate at which a

given enzyme yields one-half its maximum velocity in an enzyme catalyzed reaction.

### **I. Detection Of Specific Nucleic Acid Sequences Using 5' Nucleases**

5 The 5' nucleases of the invention form the basis of a novel detection assay for the identification of specific nucleic acid sequences. This detection system identifies the presence of specific nucleic acid sequences by requiring the annealing of two oligonucleotide probes to two portions of the target sequence. As used herein, the term "target sequence" or target nucleic acid sequence" refers to a specific nucleic acid sequence within a polynucleotide sequence, such as genomic  
10 DNA or RNA, which is to be either detected or cleaved or both.

Figure 1A provides a schematic of one embodiment of the detection method of the present invention. The target sequence is recognized by two distinct oligonucleotides in the triggering or trigger reaction. It is preferred that one of these oligonucleotides is provided on a solid support. The other can be provided  
15 free. In Figure 1A the free oligo is indicated as a "primer" and the other oligo is shown attached to a bead designated as type 1. The target nucleic acid aligns the two oligonucleotides for specific cleavage of the 5' arm (of the oligo on bead 1) by the DNAPs of the present invention (not shown in Figure 1A).

The site of cleavage (indicated by a large solid arrowhead) is controlled by  
20 the distance between the 3' end of the "primer" and the downstream fork of the oligo on bead 1. The latter is designed with an uncleavable region (indicated by the striping). In this manner neither oligonucleotide is subject to cleavage when misaligned or when unattached to target nucleic acid.

Successful cleavage releases a single copy of what is referred to as the alpha  
25 signal oligo. This oligo may contain a detectable moiety (*e.g.*, fluorescein). On the other hand, it may be unlabelled.

In one embodiment of the detection method, two more oligonucleotides are provided on solid supports. The oligonucleotide shown in Figure 1A on bead 2 has a region that is complementary to the alpha signal oligo (indicated as alpha prime)  
30 allowing for hybridization. This structure can be cleaved by the DNAPs of the present invention to release the beta signal oligo. The beta signal oligo can then

hybridize to type 3 beads having an oligo with a complementary region (indicated as beta prime). Again, this structure can be cleaved by the DNAPs of the present invention to release a new alpha oligo.

5 At this point, the amplification has been linear. To increase the power of the method, it is desired that the alpha signal oligo hybridized to bead type 2 be liberated after release of the beta oligo so that it may go on to hybridize with other oligos on type 2 beads. Similarly, after release of an alpha oligo from type 3 beads, it is desired that the beta oligo be liberated.

10 The liberation of "captured" signal oligos can be achieved in a number of ways. First, it has been found that the DNAPs of the present invention have a true 5' exonuclease capable of "nibbling" the 5' end of the alpha (and beta) prime oligo (discussed below in more detail). Thus, under appropriate conditions, the hybridization is destabilized by nibbling of the DNAP. Second, the alpha - alpha prime (as well as the beta - beta prime) complex can be destabilized by heat (*e.g.*,  
15 thermal cycling).

With the liberation of signal oligos by such techniques, each cleavage results in a doubling of the number of signal oligos. In this manner, detectable signal can quickly be achieved.

20 Figure 1B provides a schematic of a second embodiment of the detection method of the present invention. Again, the target sequence is recognized by two distinct oligonucleotides in the triggering or trigger reaction and the target nucleic acid aligns the two oligonucleotides for specific cleavage of the 5' arm by the DNAPs of the present invention (not shown in Figure 1B). The first oligo is completely complementary to a portion of the target sequence. The second  
25 oligonucleotide is partially complementary to the target sequence; the 3' end of the second oligonucleotide is fully complementary to the target sequence while the 5' end is non-complementary and forms a single-stranded arm. The non-complementary end of the second oligonucleotide may be a generic sequence which can be used with a set of standard hairpin structures (described below). The  
30 detection of different target sequences would require unique portions of two oligonucleotides: the entire first oligonucleotide and the 3' end of the second

oligonucleotide. The 5' arm of the second oligonucleotide can be invariant or generic in sequence.

5 The annealing of the first and second oligonucleotides near one another along the target sequence forms a forked cleavage structure which is a substrate for the 5' nuclease of DNA polymerases. The approximate location of the cleavage site is again indicated by the large solid arrowhead in Figure 1B.

10 The 5' nucleases of the invention are capable of cleaving this structure but are not capable of polymerizing the extension of the 3' end of the first oligonucleotide. The lack of polymerization activity is advantageous as extension of the first oligonucleotide results in displacement of the annealed region of the second oligonucleotide and results in moving the site of cleavage along the second oligonucleotide. If polymerization is allowed to occur to any significant amount, multiple lengths of cleavage product will be generated. A single cleavage product of uniform length is desirable as this cleavage product initiates the detection  
15 reaction.

The trigger reaction may be run under conditions that allow for thermocycling. Thermocycling of the reaction allows for a logarithmic increase in the amount of the trigger oligonucleotide released in the reaction.

20 The second part of the detection method allows the annealing of the fragment of the second oligonucleotide liberated by the cleavage of the first cleavage structure formed in the triggering reaction (called the third or trigger oligonucleotide) to a first hairpin structure. This first hairpin structure has a single-stranded 5' arm and a single-stranded 3' arm. The third oligonucleotide triggers the cleavage of this first hairpin structure by annealing to the 3' arm of the hairpin  
25 thereby forming a substrate for cleavage by the 5' nuclease of the present invention. The cleavage of this first hairpin structure generates two reaction products: 1) the cleaved 5' arm of the hairpin called the fourth oligonucleotide, and 2) the cleaved hairpin structure which now lacks the 5' arm and is smaller in size than the uncleaved hairpin. This cleaved first hairpin may be used as a  
30 detection molecule to indicate that cleavage directed by the trigger or third oligonucleotide occurred. Thus, this indicates that the first two oligonucleotides



found and annealed to the target sequence thereby indicating the presence of the target sequence in the sample.

5 The detection products are amplified by having the fourth oligonucleotide anneal to a second hairpin structure. This hairpin structure has a 5' single-stranded arm and a 3' single-stranded arm. The fourth oligonucleotide generated by cleavage of the first hairpin structure anneals to the 3' arm of the second hairpin structure thereby creating a third cleavage structure recognized by the 5' nuclease. The cleavage of this second hairpin structure also generates two reaction products: 1) the cleaved 5' arm of the hairpin called the fifth oligonucleotide which is similar or identical in sequence to the third nucleotide, and 2) the cleaved second hairpin structure which now lacks the 5' arm and is smaller in size than the uncleaved hairpin. This cleaved second hairpin may be as a detection molecule and amplifies the signal generated by the cleavage of the first hairpin structure. Simultaneously with the annealing of the forth oligonucleotide, the third oligonucleotide is  
10 dissociated from the cleaved first hairpin molecule so that it is free to anneal to a new copy of the first hairpin structure. The disassociation of the oligonucleotides from the hairpin structures may be accomplished by heating or other means suitable to disrupt base-pairing interactions.  
15

Further amplification of the detection signal is achieved by annealing the fifth oligonucleotide (similar or identical in sequence to the third oligonucleotide) to another molecule of the first hairpin structure. Cleavage is then performed and the oligonucleotide that is liberated then is annealed to another molecule of the second hairpin structure. Successive rounds of annealing and cleavage of the first and second hairpin structures, provided in excess, are performed to generate a  
20 sufficient amount of cleaved hairpin products to be detected. The temperature of the detection reaction is cycled just below and just above the annealing temperature for the oligonucleotides used to direct cleavage of the hairpin structures, generally about 55°C to 70°C. The number of cleavages will double in each cycle until the amount of hairpin structures remaining is below the  $K_m$  for the hairpin structures.  
25 This point is reached when the hairpin structures are substantially used up. When the detection reaction is to be used in a quantitative manner, the cycling reactions  
30

are stopped before the accumulation of the cleaved hairpin detection products reach a plateau.

Detection of the cleaved hairpin structures may be achieved in several ways. In one embodiment detection is achieved by separation on agarose or  
5 polyacrylamide gels followed by staining with ethidium bromide. In another embodiment, detection is achieved by separation of the cleaved and uncleaved hairpin structures on a gel followed by autoradiography when the hairpin structures are first labelled with a radioactive probe and separation on chromatography columns using HPLC or FPLC followed by detection of the differently sized  
10 fragments by absorption at  $OD_{260}$ . Other means of detection include detection of changes in fluorescence polarization when the single-stranded 5' arm is released by cleavage, the increase in fluorescence of an intercalating fluorescent indicator as the amount of primers annealed to 3' arms of the hairpin structures increases. The formation of increasing amounts of duplex DNA (between the primer and the 3'  
15 arm of the hairpin) occurs if successive rounds of cleavage occur.

The hairpin structures may be attached to a solid support, such as an agarose, styrene or magnetic bead, via the 3' end of the hairpin. A spacer molecule may be placed between the 3' end of the hairpin and the bead, if so desired. The advantage of attaching the hairpin structures to a solid support is that  
20 this prevents the hybridization of the two hairpin structures to one another over regions which are complementary. If the hairpin structures anneal to one another, this would reduce the amount of hairpins available for hybridization to the primers released during the cleavage reactions. If the hairpin structures are attached to a solid support, then additional methods of detection of the products of the cleavage  
25 reaction may be employed. These methods include, but are not limited to, the measurement of the released single-stranded 5' arm when the 5' arm contains a label at the 5' terminus. This label may be radioactive, fluorescent, biotinylated, etc. If the hairpin structure is not cleaved, the 5' label will remain attached to the solid support. If cleavage occurs, the 5' label will be released from the solid  
30 support.

The 3' end of the hairpin molecule may be blocked through the use of dideoxynucleotides. A 3' terminus containing a dideoxynucleotide is unavailable to



participate in reactions with certain DNA modifying enzymes, such as terminal transferase. Cleavage of the hairpin having a 3' terminal dideoxynucleotide generates a new, unblocked 3' terminus at the site of cleavage. This new 3' end has a free hydroxyl group which can interact with terminal transferase thus providing another means of detecting the cleavage products.

The hairpin structures are designed so that their self-complementary regions are very short (generally in the range of 3-8 base pairs). Thus, the hairpin structures are not stable at the high temperatures at which this reaction is performed (generally in the range of 50-75°C) unless the hairpin is stabilized by the presence of the annealed oligonucleotide on the 3' arm of the hairpin. This instability prevents the polymerase from cleaving the hairpin structure in the absence of an associated primer thereby preventing false positive results due to non-oligonucleotide directed cleavage.

As discussed above, the use of the 5' nucleases of the invention which have reduced polymerization activity is advantageous in this method of detecting specific nucleic acid sequences. Significant amounts of polymerization during the cleavage reaction would cause shifting of the site of cleavage in unpredictable ways resulting in the production of a series of cleaved hairpin structures of various sizes rather than a single easily quantifiable product. Additionally, the primers used in one round of cleavage could, if elongated, become unusable for the next cycle, by either forming an incorrect structure or by being too long to melt off under moderate temperature cycling conditions. In a pristine system (*i.e.*, lacking the presence of dNTPs), one could use the unmodified polymerase, but the presence of nucleotides (dNTPs) can decrease the per cycle efficiency enough to give a false negative result. When a crude extract (genomic DNA preparations, crude cell lysates, *etc.*) is employed or where a sample of DNA from a PCR reaction, or any other sample that might be contaminated with dNTPs, the 5' nucleases of the present invention that were derived from thermostable polymerases are particularly useful.

## II. Generation Of 5' Nucleases From Thermostable DNA Polymerases

The genes encoding Type A DNA polymerases share about 85% homology to each other on the DNA sequence level. Preferred examples of thermostable polymerases include those isolated from *Thermus aquaticus*, *Thermus flavus*, and *Thermus thermophilus*. However, other thermostable Type A polymerases which have 5' nuclease activity are also suitable. Figs. 2 and 3 compare the nucleotide and amino acid sequences of the three above mentioned polymerases. SEQ ID NOS:1-3 display the nucleotide sequences and SEQ ID NOS:4-6 display the amino acid sequences of the three wild-type polymerases. SEQ ID NO:1 corresponds to the nucleic acid sequence of the wild type *Thermus aquaticus* DNA polymerase gene isolated from the YT-1 strain [Lawyer *et al.*, *J. Biol. Chem.* 264:6427 (1989)]. SEQ ID NO:2 corresponds to the nucleic acid sequence of the wild type *Thermus flavus* DNA polymerase gene [Akhmetzjanov and Vakhitov, *Nucl. Acids Res.* 20:5839 (1992)]. SEQ ID NO:3 corresponds to the nucleic acid sequence of the wild type *Thermus thermophilus* DNA polymerase gene [Gelfand *et al.*, WO 91/09950 (1991)]. SEQ ID NOS:7-8 depict the consensus nucleotide and amino acid sequences, respectively for the above three DNAPs (also shown on the top row in Figs. 2 and 3).

The 5' nucleases of the invention derived from thermostable polymerases have reduced synthetic ability, but retain substantially the same 5' exonuclease activity as the native DNA polymerase. The term "substantially the same 5' nuclease activity" as used herein means that the 5' nuclease activity of the modified enzyme retains the ability to function as a structure-dependent single-stranded endonuclease but not necessarily at the same rate of cleavage as compared to the unmodified enzyme. Type A DNA polymerases may also be modified so as to produce an enzyme which has increased 5' nuclease activity while having a reduced level of synthetic activity. Modified enzymes having reduced synthetic activity and increased 5' nuclease activity are also envisioned by the present invention.

By the term "reduced synthetic activity" as used herein it is meant that the modified enzyme has less than the level of synthetic activity found in the unmodified or "native" enzyme. The modified enzyme may have no synthetic

activity remaining or may have that level of synthetic activity that will not interfere with the use of the modified enzyme in the detection assay described below. The 5' nucleases of the present invention are advantageous in situations where the cleavage activity of the polymerase is desired, but the synthetic ability is not (such as in the detection assay of the invention).

As noted above, it is not intended that the invention be limited by the nature of the alteration necessary to render the polymerase synthesis deficient. The present invention contemplates a variety of methods, including but not limited to: 1) proteolysis; 2) recombinant constructs (including mutants); and 3) physical and/or chemical modification and/or inhibition.

### 1. Proteolysis

Thermostable DNA polymerases having a reduced level of synthetic activity are produced by physically cleaving the unmodified enzyme with proteolytic enzymes to produce fragments of the enzyme that are deficient in synthetic activity but retain 5' nuclease activity. Following proteolytic digestion, the resulting fragments are separated by standard chromatographic techniques and assayed for the ability to synthesize DNA and to act as a 5' nuclease. The assays to determine synthetic activity and 5' nuclease activity are described below.

### 2. Recombinant Constructs

The examples below describe a preferred method for creating a construct encoding a 5' nuclease derived from a thermostable DNA polymerase. As the Type A DNA polymerases are similar in DNA sequence, the cloning strategies employed for the *Thermus aquaticus* and *flavus* polymerases are applicable to other thermostable Type A polymerases. In general, a thermostable DNA polymerase is cloned by isolating genomic DNA using molecular biological methods from a bacteria containing a thermostable Type A DNA polymerase. This genomic DNA is exposed to primers which are capable of amplifying the polymerase gene by PCR.

This amplified polymerase sequence is then subjected to standard deletion processes to delete the polymerase portion of the gene. Suitable deletion processes are described below in the examples.

5 The example below discusses the strategy used to determine which portions of the DNAP*Taq* polymerase domain could be removed without eliminating the 5' nuclease activity. Deletion of amino acids from the protein can be done either by deletion of the encoding genetic material, or by introduction of a translational stop codon by mutation or frame shift. In addition, proteolytic treatment of the protein molecule can be performed to remove segments of the protein.

10 In the examples below, specific alterations of the *Taq* gene were: a deletion between nucleotides 1601 and 2502 (the end of the coding region), a 4 nucleotide insertion at position 2043, and deletions between nucleotides 1614 and 1848 and between nucleotides 875 and 1778 (numbering is as in SEQ ID NO:1). These modified sequences are described below in the examples and at SEQ ID NOS:9-12.

15 Those skilled in the art understand that single base pair changes can be innocuous in terms of enzyme structure and function. Similarly, small additions and deletions can be present without substantially changing the exonuclease or polymerase function of these enzymes.

20 Other deletions are also suitable to create the 5' nucleases of the present invention. It is preferable that the deletion decrease the polymerase activity of the 5' nucleases to a level at which synthetic activity will not interfere with the use of the 5' nuclease in the detection assay of the invention. Most preferably, the synthetic ability is absent. Modified polymerases are tested for the presence of synthetic and 5' nuclease activity as in assays described below. Thoughtful  
25 consideration of these assays allows for the screening of candidate enzymes whose structure is heretofore as yet unknown. In other words, construct "X" can be evaluated according to the protocol described below to determine whether it is a member of the genus of 5' nucleases of the present invention as defined functionally, rather than structurally.

30 In the example below, the PCR product of the amplified *Thermus aquaticus* genomic DNA did not have the identical nucleotide structure of the native genomic DNA and did not have the same synthetic ability of the original clone. Base pair

changes which result due to the infidelity of DNAP $Taq$  during PCR amplification of a polymerase gene are also a method by which the synthetic ability of a polymerase gene may be inactivated. The examples below and Figs. 4A and 5A indicate regions in the native *Thermus aquaticus* and *flavus* DNA polymerases likely to be important for synthetic ability. There are other base pair changes and substitutions that will likely also inactivate the polymerase.

It is not necessary, however, that one start out the process of producing a 5' nuclease from a DNA polymerase with such a mutated amplified product. This is the method by which the examples below were performed to generate the synthesis-deficient DNAP $Taq$  mutants, but it is understood by those skilled in the art that a wild-type DNA polymerase sequence may be used as the starting material for the introduction of deletions, insertion and substitutions to produce a 5' nuclease. For example, to generate the synthesis-deficient DNAPT $fl$  mutant, the primers listed in SEQ ID NOS:13-14 were used to amplify the wild type DNA polymerase gene from *Thermus flavus* strain AT-62. The amplified polymerase gene was then subjected to restriction enzyme digestion to delete a large portion of the domain encoding the synthetic activity.

The present invention contemplates that the nucleic acid construct of the present invention be capable of expression in a suitable host. Those in the art know methods for attaching various promoters and 3' sequences to a gene structure to achieve efficient expression. The examples below disclose two suitable vectors and six suitable vector constructs. Of course, there are other promoter/vector combinations that would be suitable. It is not necessary that a host organism be used for the expression of the nucleic acid constructs of the invention. For example, expression of the protein encoded by a nucleic acid construct may be achieved through the use of a cell-free in vitro transcription/translation system. An example of such a cell-free system is the commercially available TnT<sup>TM</sup> Coupled Reticulocyte Lysate System (Promega Corporation, Madison, WI).

Once a suitable nucleic acid construct has been made, the 5' nuclease may be produced from the construct. The examples below and standard molecular biological teachings enable one to manipulate the construct by different suitable methods.

Once the 5' nuclease has been expressed, the polymerase is tested for both synthetic and nuclease activity as described below.

### 3. Physical And/Or Chemical Modification And/Or Inhibition

5           The synthetic activity of a thermostable DNA polymerase may be reduced by chemical and/or physical means. In one embodiment, the cleavage reaction catalyzed by the 5' nuclease activity of the polymerase is run under conditions which preferentially inhibit the synthetic activity of the polymerase. The level of synthetic activity need only be reduced to that level of activity which does not  
10           interfere with cleavage reactions requiring no significant synthetic activity.

          As shown in the examples below, concentrations of  $Mg^{++}$  greater than 5 mM inhibit the polymerization activity of the native DNAPTaq. The ability of the 5' nuclease to function under conditions where synthetic activity is inhibited is tested by running the assays for synthetic and 5' nuclease activity, described below, in the  
15           presence of a range of  $Mg^{++}$  concentrations (5 to 10 mM). The effect of a given concentration of  $Mg^{++}$  is determined by quantitation of the amount of synthesis and cleavage in the test reaction as compared to the standard reaction for each assay.

          The inhibitory effect of other ions, polyamines, denaturants, such as urea, formamide, dimethylsulfoxide, glycerol and non-ionic detergents (Triton X-100 and  
20           Tween-20), nucleic acid binding chemicals such as, actinomycin D, ethidium bromide and psoralens, are tested by their addition to the standard reaction buffers for the synthesis and 5' nuclease assays. Those compounds having a preferential inhibitory effect on the synthetic activity of a thermostable polymerase are then used to create reaction conditions under which 5' nuclease activity (cleavage) is  
25           retained while synthetic activity is reduced or eliminated.

          Physical means may be used to preferentially inhibit the synthetic activity of a polymerase. For example, the synthetic activity of thermostable polymerases is destroyed by exposure of the polymerase to extreme heat (typically 96 to 100°C) for extended periods of time (greater than or equal to 20 minutes). While these are  
30           minor differences with respect to the specific heat tolerance for each of the enzymes, these are readily determined. Polymerases are treated with heat for



various periods of time and the effect of the heat treatment upon the synthetic and 5' nuclease activities is determined.

### III. Therapeutic Utility Of 5' Nucleases

5 The 5' nucleases of the invention have not only the diagnostic utility discussed above, but additionally have therapeutic utility for the cleavage and inactivation of specific mRNAs inside infected cells. The mRNAs of pathogenic agents, such as viruses, bacteria, are targeted for cleavage by a synthesis-deficient DNA polymerase by the introduction of an oligonucleotide complementary to a given mRNA produced by the pathogenic agent into the infected cell along with the  
10 synthesis-deficient polymerase. Any pathogenic agent may be targeted by this method provided the nucleotide sequence information is available so that an appropriate oligonucleotide may be synthesized. The synthetic oligonucleotide anneals to the complementary mRNA thereby forming a cleavage structure recognized by the modified enzyme. The ability of the 5' nuclease activity of  
15 thermostable DNA polymerases to cleave RNA-DNA hybrids is shown herein in Example 1D.

Liposomes provide a convenient delivery system. The synthetic oligonucleotide may be conjugated or bound to the nuclease to allow for co-delivery of these molecules. Additional delivery systems may be employed.

20 Inactivation of pathogenic mRNAs has been described using antisense gene regulation and using ribozymes (Rossi, U.S. Patent No. 5,144,019, hereby incorporated by reference). Both of these methodologies have limitations.

The use of antisense RNA to impair gene expression requires stoichiometric and therefore, large molar excesses of anti-sense RNA relative to the pathogenic  
25 RNA to be effective. Ribozyme therapy, on the other hand, is catalytic and therefore lacks the problem of the need for a large molar excess of the therapeutic compound found with antisense methods. However, ribozyme cleavage of a given RNA requires the presence of highly conserved sequences to form the catalytically active cleavage structure. This requires that the target pathogenic mRNA contain  
30 the conserved sequences (GAAAC (X)<sub>n</sub> GU) thereby limiting the number of pathogenic mRNAs that can be cleaved by this method. In contrast, the catalytic

cleavage of RNA by the use of a DNA oligonucleotide and a 5' nuclease is dependent upon structure only; thus, virtually any pathogenic RNA sequence can be used to design an appropriate cleavage structure.

#### IV. Detection Of Antigenic Or Nucleic Acid Targets By A Dual Capture Assay

The ability to generate 5' nucleases from thermostable DNA polymerases provides the basis for a novel means of detecting the presence of antigenic or nucleic acid targets. In this dual capture assay, the polymerase domains encoding the synthetic activity and the nuclease activity are covalently attached to two separate and distinct antibodies or oligonucleotides. When both the synthetic and the nuclease domains are present in the same reaction and dATP, dTTP and a small amount of poly d(A-T) are provided, an enormous amount of poly d(A-T) is produced. The large amounts of poly d(A-T) are produced as a result of the ability of the 5' nuclease to cleave newly made poly d(A-T) to generate primers that are, in turn, used by the synthetic domain to catalyze the production of even more poly d(A-T). The 5' nuclease is able to cleave poly d(A-T) because poly d(A-T) is self-complementary and easily forms alternate structures at elevated temperatures. These structures are recognized by the 5' nuclease and are then cleaved to generate more primer for the synthesis reaction.

The following is an example of the dual capture assay to detect an antigen(s): A sample to be analyzed for a given antigen(s) is provided. This sample may comprise a mixture of cells; for example, cells infected with viruses display virally-encoded antigens on their surface. If the antigen(s) to be detected are present in solution, they are first attached to a solid support such as the wall of a microtiter dish or to a bead using conventional methodologies. The sample is then mixed with 1) the synthetic domain of a thermostable DNA polymerase conjugated to an antibody which recognizes either a first antigen or a first epitope on an antigen, and 2) the 5' nuclease domain of a thermostable DNA polymerase conjugated to a second antibody which recognizes either a second, distinct antigen or a second epitope on the same antigen as recognized by the antibody conjugated to the synthetic domain. Following an appropriate period to allow the interaction



of the antibodies with their cognate antigens (conditions will vary depending upon the antibodies used; appropriate conditions are well known in the art), the sample is then washed to remove unbound antibody-enzyme domain complexes. dATP, dTTP and a small amount of poly d(A-T) is then added to the washed sample and the sample is incubated at elevated temperatures (generally in the range of 60-80°C and more preferably, 70-75°C) to permit the thermostable synthetic and 5' nuclease domains to function. If the sample contains the antigen(s) recognized by both separately conjugated domains of the polymerase, then an exponential increase in poly d(A-T) production occurs. If only the antibody conjugated to the synthetic domain of the polymerase is present in the sample such that no 5' nuclease domain is present in the washed sample, then only an arithmetic increase in poly d(A-T) is possible. The reaction conditions may be controlled in such a way so that an arithmetic increase in poly d(A-T) is below the threshold of detection. This may be accomplished by controlling the length of time the reaction is allowed to proceed or by adding so little poly d(A-T) to act as template that in the absence of nuclease activity to generate new poly d(A-T) primers very little poly d(A-T) is synthesized.

It is not necessary for both domains of the enzyme to be conjugated to an antibody. One can provide the synthetic domain conjugated to an antibody and provide the 5' nuclease domain in solution or vice versa. In such a case the conjugated antibody-enzyme domain is added to the sample, incubated, then washed. dATP, dTTP, poly d(A-T) and the remaining enzyme domain in solution is then added.

Additionally, the two enzyme domains may be conjugated to oligonucleotides such that target nucleic acid sequences can be detected. The oligonucleotides conjugated to the two different enzyme domains may recognize different regions on the same target nucleic acid strand or may recognize two unrelated target nucleic acids.

The production of poly d(A-T) may be detected in many ways including:  
1) use of a radioactive label on either the dATP or dTTP supplied for the synthesis of the poly d(A-T), followed by size separation of the reaction products and autoradiography; 2) use of a fluorescent probe on the dATP and a biotinylated

probe on the dTTP supplied for the synthesis of the poly d(A-T), followed by passage of the reaction products over an avidin bead, such as magnetic beads conjugated to avidin; the presence of the florescent probe on the avidin-containing bead indicates that poly d(A-T) has been formed as the fluorescent probe will stick to the avidin bead only if the fluorescenated dATP is incorporated into a covalent linkage with the biotinylated dTTP; and 3) changes fluorescence polarization indicating an increase in size. Other means of detecting the presence of poly d(A-T) include the use of intercalating fluorescence indicators to monitor the increase in duplex DNA formation.

The advantages of the above dual capture assay for detecting antigenic or nucleic acid targets include:

1) No thermocycling of the sample is required. The polymerase domains and the dATP and dTTP are incubated at a fixed temperature (generally about 70°C). After 30 minutes of incubation up to 75% of the added dNTPs are incorporated into poly d(A-T). The lack of thermocycling makes this assay well suited to clinical laboratory settings; there is no need to purchase a thermocycling apparatus and there is no need to maintain very precise temperature control.

2) The reaction conditions are simple. The incubation of the bound enzymatic domains is done in a buffer containing 0.5 mM MgCl<sub>2</sub> (higher concentrations may be used), 2-10 mM Tris-Cl, pH 8.5, approximately 50 μM dATP and dTTP. The reaction volume is 10-20 μl and reaction products are detectable within 10-20 minutes.

3) No reaction is detected unless both the synthetic and nuclease activities are present. Thus, a positive result indicates that both probes (antibody or oligonucleotide) have recognized their targets thereby increasing the specificity of recognition by having two different probes bind to the target.

The ability to separate the two enzymatic activities of the DNAP allows for exponential increases in poly d(A-T) production. If a DNAP is used which lacks 5' nuclease activity, such as the Klenow fragment of DNAPEc1, only a linear or arithmetic increase in poly d(A-T) production is possible [Setlow *et al.*, J. Biol. Chem. 247:224 (1972)]. The ability to provide an enzyme having 5' nuclease

activity but lacking synthetic activity is made possible by the disclosure of this invention.

### EXPERIMENTAL

5 The following examples serve to illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

In the disclosure which follows, the following abbreviations apply: °C (degrees Centigrade); g (gravitational field); vol (volume); w/v (weight to volume); v/v (volume to volume); BSA (bovine serum albumin); CTAB  
10 (cetyltrimethylammonium bromide); HPLC (high pressure liquid chromatography); DNA (deoxyribonucleic acid); p (plasmid); µl (microliters); ml (milliliters); µg (micrograms); pmoles (picomoles); mg (milligrams); M (molar); mM (milliMolar); µM (microMolar); nm (nanometers); kdal (kilodaltons); OD (optical density); EDTA (ethylene diamine tetra-acetic acid); FITC (fluorescein isothiocyanate); SDS  
15 (sodium dodecyl sulfate); NaPO<sub>4</sub> (sodium phosphate); Tris (tris(hydroxymethyl)-aminomethane); PMSF (phenylmethylsulfonylfluoride); TBE (Tris-Borate-EDTA, *i.e.*, Tris buffer titrated with boric acid rather than HCl and containing EDTA) ; PBS (phosphate buffered saline); PPBS (phosphate buffered saline containing 1 mM PMSF); PAGE (polyacrylamide gel electrophoresis); Tween (polyoxyethylene-sorbitan); Dynal (Dynal A.S., Oslo, Norway); Epicentre (Epicentre Technologies,  
20 Madison, WI); National Biosciences (Plymouth, MN); New England Biolabs (Beverly, MA); Novagen (Novagen, Inc., Madison, WI); Perkin Elmer (Norwalk, CT); Promega Corp. (Madison, WI); Stratagene (Stratagene Cloning Systems, La Jolla, CA); USB (U.S. Biochemical, Cleveland, OH).

## EXAMPLE 1

## Characteristics Of Native Thermostable DNA Polymerases

A. 5' Nuclease Activity Of DNAP*Taq*

During the polymerase chain reaction (PCR) [Saiki *et al.*, *Science* 239:487  
5 (1988); Mullis and Faloona, *Methods in Enzymology* 155:335 (1987)], DNAP*Taq* is  
able to amplify many, but not all, DNA sequences. One sequence that cannot be  
amplified using DNAP*Taq* is shown in Figure 6 (Hairpin structure is SEQ ID  
NO:15, PRIMERS are SEQ ID NOS:16-17.) This DNA sequence has the  
distinguishing characteristic of being able to fold on itself to form a hairpin with  
10 two single-stranded arms, which correspond to the primers used in PCR.

To test whether this failure to amplify is due to the 5' nuclease activity of  
the enzyme, we compared the abilities of DNAP*Taq* and DNAPStf to amplify this  
DNA sequence during 30 cycles of PCR. Synthetic oligonucleotides were obtained  
from The Biotechnology Center at the University of Wisconsin-Madison. The  
15 DNAP*Taq* and DNAPStf were from Perkin Elmer (*i.e.*, Amplitaq™ DNA  
polymerase and the Stoffel fragment of Amplitaq™ DNA polymerase). The  
substrate DNA comprised the hairpin structure shown in Figure 6 cloned in a  
double-stranded form into pUC19. The primers used in the amplification are listed  
as SEQ ID NOS:16-17. Primer SEQ ID NO:17 is shown annealed to the 3' arm of  
20 the hairpin structure in Fig. 6. Primer SEQ ID NO:16 is shown as the first 20  
nucleotides in bold on the 5' arm of the hairpin in Fig. 6.

Polymerase chain reactions comprised 1 ng of supercoiled plasmid target  
DNA, 5 pmoles of each primer, 40 μM each dNTP, and 2.5 units of DNAP*Taq* or  
DNAPStf, in a 50 μl solution of 10 mM Tris•Cl pH 8.3. The DNAP*Taq* reactions  
25 included 50 mM KCl and 1.5 mM MgCl<sub>2</sub>. The temperature profile was 95°C for  
30 sec., 55°C for 1 min. and 72°C for 1 min., through 30 cycles. Ten percent of  
each reaction was analyzed by gel electrophoresis through 6% polyacrylamide  
(cross-linked 29:1) in a buffer of 45 mM Tris•Borate, pH 8.3, 1.4 mM EDTA.

The results are shown in Figure 7. The expected product was made by  
30 DNAPStf (indicated simply as "S") but not by DNAP*Taq* (indicated as "T"). We

conclude that the 5' nuclease activity of DNAP*Taq* is responsible for the lack of amplification of this DNA sequence.

To test whether the 5' unpaired nucleotides in the substrate region of this structured DNA are removed by DNAP*Taq*, the fate of the end-labeled 5' arm during four cycles of PCR was compared using the same two polymerases (Figure. 8). The hairpin templates, such as the one described in Figure 6, were made using DNAP*Stf* and a <sup>32</sup>P-5'-end-labeled primer. The 5'-end of the DNA was released as a few large fragments by DNAP*Taq* but not by DNAP*Stf*. The sizes of these fragments (based on their mobilities) show that they contain most or all of the unpaired 5' arm of the DNA. Thus, cleavage occurs at or near the base of the bifurcated duplex. These released fragments terminate with 3' OH groups, as evidenced by direct sequence analysis, and the abilities of the fragments to be extended by terminal deoxynucleotidyl transferase.

Figures 9-11 show the results of experiments designed to characterize the cleavage reaction catalyzed by DNAP*Taq*. Unless otherwise specified, the cleavage reactions comprised 0.01 pmoles of heat-denatured, end-labeled hairpin DNA (with the unlabeled complementary strand also present), 1 pmole primer (complementary to the 3' arm) and 0.5 units of DNAP*Taq* (estimated to be 0.026 pmoles) in a total volume of 10  $\mu$ l of 10 mM Tris-Cl, pH 8.5, 50 mM KCl and 1.5 mM MgCl<sub>2</sub>. As indicated, some reactions had different concentrations of KCl, and the precise times and temperatures used in each experiment are indicated in the individual figures. The reactions that included a primer used the one shown in Figure 6 (SEQ ID NO:17). In some instances, the primer was extended to the junction site by providing polymerase and selected nucleotides.

Reactions were initiated at the final reaction temperature by the addition of either the MgCl<sub>2</sub> or enzyme. Reactions were stopped at their incubation temperatures by the addition of 8  $\mu$ l of 95% formamide with 20 mM EDTA and 0.05% marker dyes. The  $T_m$  calculations listed were made using the Oligo<sup>TM</sup> primer analysis software from National Biosciences, Inc. These were determined using 0.25  $\mu$ M as the DNA concentration, at either 15 or 65 mM total salt (the 1.5 mM MgCl<sub>2</sub> in all reactions was given the value of 15 mM salt for these calculations).

Figure 9 is an autoradiogram containing the results of a set of experiments and conditions on the cleavage site. Figure 9A is a determination of reaction components that enable cleavage. Incubation of 5'-end-labeled hairpin DNA was for 30 minutes at 55°C, with the indicated components. The products were resolved by denaturing polyacrylamide gel electrophoresis and the lengths of the products, in nucleotides, are indicated. Figure 9B describes the effect of temperature on the site of cleavage in the absence of added primer. Reactions were incubated in the absence of KCl for 10 minutes at the indicated temperatures. The lengths of the products, in nucleotides, are indicated.

Surprisingly, cleavage by DNAP*Taq* requires neither a primer nor dNTPs (see Fig. 9A). Thus, the 5' nuclease activity can be uncoupled from polymerization. Nuclease activity requires magnesium ions, though manganese ions can be substituted, albeit with potential changes in specificity and activity. Neither zinc nor calcium ions support the cleavage reaction. The reaction occurs over a broad temperature range, from 25°C to 85°C, with the rate of cleavage increasing at higher temperatures.

Still referring to Figure 9, the primer is not elongated in the absence of added dNTPs. However, the primer influences both the site and the rate of cleavage of the hairpin. The change in the site of cleavage (Fig. 9A) apparently results from disruption of a short duplex formed between the arms of the DNA substrate. In the absence of primer, the sequences indicated by underlining in Figure 6 could pair, forming an extended duplex. Cleavage at the end of the extended duplex would release the 11 nucleotide fragment seen on the Fig. 9A lanes with no added primer. Addition of excess primer (Fig. 9A, lanes 3 and 4) or incubation at an elevated temperature (Fig. 9B) disrupts the short extension of the duplex and results in a longer 5' arm and, hence, longer cleavage products.

The location of the 3' end of the primer can influence the precise site of cleavage. Electrophoretic analysis revealed that in the absence of primer (Fig. 9B), cleavage occurs at the end of the substrate duplex (either the extended or shortened form, depending on the temperature) between the first and second base pairs. When the primer extends up to the base of the duplex, cleavage also occurs one nucleotide into the duplex. However, when a gap of four or six nucleotides exists



between the 3' end of the primer and the substrate duplex, the cleavage site is shifted four to six nucleotides in the 5' direction.

Fig. 10 describes the kinetics of cleavage in the presence (Fig. 10A) or absence (Fig. 10B) of a primer oligonucleotide. The reactions were run at 55°C with either 50 mM KCl (Fig. 10A) or 20 mM KCl (Fig. 10B). The reaction products were resolved by denaturing polyacrylamide gel electrophoresis and the lengths of the products, in nucleotides, are indicated. "M", indicating a marker, is a 5' end-labeled 19-nt oligonucleotide. Under these salt conditions, Figs. 10A and 10B indicate that the reaction appears to be about twenty times faster in the presence of primer than in the absence of primer. This effect on the efficiency may be attributable to proper alignment and stabilization of the enzyme on the substrate.

The relative influence of primer on cleavage rates becomes much greater when both reactions are run in 50 mM KCl. In the presence of primer, the rate of cleavage increases with KCl concentration, up to about 50 mM. However, inhibition of this reaction in the presence of primer is apparent at 100 mM and is complete at 150 mM KCl. In contrast, in the absence of primer the rate is enhanced by concentration of KCl up to 20 mM, but it is reduced at concentrations above 30 mM. At 50 mM KCl, the reaction is almost completely inhibited. The inhibition of cleavage by KCl in the absence of primer is affected by temperature, being more pronounced at lower temperatures.

Recognition of the 5' end of the arm to be cut appears to be an important feature of substrate recognition. Substrates that lack a free 5' end, such as circular M13 DNA, cannot be cleaved under any conditions tested. Even with substrates having defined 5' arms, the rate of cleavage by DNAP*Taq* is influenced by the length of the arm. In the presence of primer and 50 mM KCl, cleavage of a 5' extension that is 27 nucleotides long is essentially complete within 2 minutes at 55°C. In contrast, cleavages of molecules with 5' arms of 84 and 188 nucleotides are only about 90% and 40% complete after 20 minutes. Incubation at higher temperatures reduces the inhibitory effects of long extensions indicating that secondary structure in the 5' arm or a heat-labile structure in the enzyme may inhibit the reaction. A mixing experiment, run under conditions of substrate

excess, shows that the molecules with long arms do not preferentially tie up the available enzyme in non-productive complexes. These results may indicate that the 5' nuclease domain gains access to the cleavage site at the end of the bifurcated duplex by moving down the 5' arm from one end to the other. Longer 5' arms would be expected to have more adventitious secondary structures (particularly when KCl concentrations are high), which would be likely to impede this movement.

Cleavage does not appear to be inhibited by long 3' arms of either the substrate strand target molecule or pilot nucleic acid, at least up to 2 kilobases. At the other extreme, 3' arms of the pilot nucleic acid as short as one nucleotide can support cleavage in a primer-independent reaction, albeit inefficiently. Fully paired oligonucleotides do not elicit cleavage of DNA templates during primer extension.

The ability of DNAP $Taq$  to cleave molecules even when the complementary strand contains only one unpaired 3' nucleotide may be useful in optimizing allele-specific PCR. PCR primers that have unpaired 3' ends could act as pilot oligonucleotides to direct selective cleavage of unwanted templates during preincubation of potential template-primer complexes with DNAP $Taq$  in the absence of nucleoside triphosphates.

#### **B. 5' Nuclease Activities Of Other DNAPs**

To determine whether other 5' nucleases in other DNAPs would be suitable for the present invention, an array of enzymes, several of which were reported in the literature to be free of apparent 5' nuclease activity, were examined. The ability of these other enzymes to cleave nucleic acids in a structure-specific manner was tested using the hairpin substrate shown in Fig. 6 under conditions reported to be optimal for synthesis by each enzyme.

DNAPEcl and DNAP Klenow were obtained from Promega Corporation; the DNAP of *Pyrococcus furiosus* ["Pfu", Bargseid *et al.*, Strategies 4:34 (1991)] was from Strategene; the DNAP of *Thermococcus litoralis* ["Tli", Vent<sup>TM</sup>(exo-), Perler *et al.*, Proc. Natl. Acad. Sci. USA 89:5577 (1992)] was from New England Biolabs; the DNAP of *Thermus flavus* ["Tfl", Kaledin *et al.*, Biokhimiya 46:1576 (1981)] was from Epicentre Technologies; and the DNAP of *Thermus thermophilus*

["Tth", Carballeira *et al.*, Biotechniques 9:276 (1990); Myers *et al.*, *Biochem.* 30:7661 (1991)] was from U.S. Biochemicals.

0.5 units of each DNA polymerase was assayed in a 20  $\mu$ l reaction, using either the buffers supplied by the manufacturers for the primer-dependent reactions, or 10 mM Tris•Cl, pH 8.5, 1.5 mM MgCl<sub>2</sub>, and 20mM KCl. Reaction mixtures were at held 72°C before the addition of enzyme.

Fig. 11 is an autoradiogram recording the results of these tests. Fig. 11A demonstrates reactions of endonucleases of DNAPs of several thermophilic bacteria. The reactions were incubated at 55°C for 10 minutes in the presence of primer or at 72°C for 30 minutes in the absence of primer, and the products were resolved by denaturing polyacrylamide gel electrophoresis. The lengths of the products, in nucleotides, are indicated. Fig. 11B demonstrates endonucleolytic cleavage by the 5' nuclease of DNAPEcl. The DNAPEcl and DNAP Klenow reactions were incubated for 5 minutes at 37°C. Note the light band of cleavage products of 25 and 11 nucleotides in the DNAPEcl lanes (made in the presence and absence of primer, respectively). Fig. 7B also demonstrates DNAPTaq reactions in the presence (+) or absence (-) of primer. These reactions were run in 50 mM and 20 mM KCl, respectively, and were incubated at 55°C for 10 minutes.

Referring to Fig. 11A, DNAPs from the eubacteria *Thermus thermophilus* and *Thermus flavus* cleave the substrate at the same place as DNAPTaq, both in the presence and absence of primer. In contrast, DNAPs from the archaebacteria *Pyrococcus furiosus* and *Thermococcus litoralis* are unable to cleave the substrates endonucleolytically. The DNAPs from *Pyrococcus furiosus* and *Thermococcus litoralis* share little sequence homology with eubacterial enzymes (Ito *et al.*, *Nucl. Acids Res.* 19:4045 (1991); Mathur *et al.*, *Nucl. Acids. Res.* 19:6952 (1991); see also Perler *et al.*). Referring to Fig. 11B, DNAPEcl also cleaves the substrate, but the resulting cleavage products are difficult to detect unless the 3' exonuclease is inhibited. The amino acid sequences of the 5' nuclease domains of DNAPEcl and DNAPTaq are about 38% homologous (Gelfand, *supra*).

The 5' nuclease domain of DNAPTaq also shares about 19% homology with the 5' exonuclease encoded by gene 6 of bacteriophage T7 [Dunn *et al.*, *J. Mol. Biol.* 166:477 (1983)]. This nuclease, which is not covalently attached to a DNAP

polymerization domain, is also able to cleave DNA endonucleolytically, at a site similar or identical to the site that is cut by the 5' nucleases described above, in the absence of added primers.

### C. Transcleavage

5           The ability of a 5' nuclease to be directed to cleave efficiently at any specific sequence was demonstrated in the following experiment. A partially complementary oligonucleotide termed a "pilot oligonucleotide" was hybridized to sequences at the desired point of cleavage. The non-complementary part of the pilot oligonucleotide provided a structure analogous to the 3' arm of the template  
10           (see Fig. 6), whereas the 5' region of the substrate strand became the 5' arm. A primer was provided by designing the 3' region of the pilot so that it would fold on itself creating a short hairpin with a stabilizing tetra-loop [Antao *et al.*, *Nucl. Acids Res.* 19:5901 (1991)]. Two pilot oligonucleotides are shown in Fig. 12A. Oligonucleotides 19-12 (SEQ ID NO:18), 30-12 (SEQ ID NO:19) and 30-0 (SEQ  
15           ID NO:40) are 31, 42 or 30 nucleotides long, respectively. However, oligonucleotides 19-12 (SEQ ID NO:18) and 34-19 (SEQ ID NO:19) have only 19 and 30 nucleotides, respectively, that are complementary to different sequences in the substrate strand. The pilot oligonucleotides are calculated to melt off their complements at about 50°C (19-12) and about 75°C (30-12). Both pilots have 12  
20           nucleotides at their 3' ends, which act as 3' arms with base-paired primers attached.

          To demonstrate that cleavage could be directed by a pilot oligonucleotide, we incubated a single-stranded target DNA with DNAP $Taq$  in the presence of two potential pilot oligonucleotides. The transcleavage reactions, where the target and pilot nucleic acids are not covalently linked, includes 0.01 pmoles of single end-  
25           labeled substrate DNA, 1 unit of DNAP $Taq$  and 5 pmoles of pilot oligonucleotide in a volume of 20  $\mu$ l of the same buffers. These components were combined during a one minute incubation at 95°C, to denature the PCR-generated double-stranded substrate DNA, and the temperatures of the reactions were then reduced to their final incubation temperatures. Oligonucleotides 30-12 and 19-12 can  
30           hybridize to regions of the substrate DNAs that are 85 and 27 nucleotides from the 5' end of the targeted strand.

Figure 21 shows the complete 206-mer sequence (SEQ ID NO:32). The 206-mer was generated by PCR. The M13/pUC 24-mer reverse sequencing (-48) primer and the M13/pUC sequencing (-47) primer from New England Biolabs (catalogue nos. 1233 and 1224 respectively) were used (50 pmoles each) with the pGEM3z(f+) plasmid vector (Promega Corp.) as template (10 ng) containing the target sequences. The conditions for PCR were as follows: 50  $\mu$ M of each dNTP and 2.5 units of Taq DNA polymerase in 100  $\mu$ l of 20 mM Tris-Cl, pH 8.3, 1.5 mM  $MgCl_2$ , 50 mM KCl with 0.05% Tween-20 and 0.05% NP-40. Reactions were cycled 35 times through 95°C for 45 seconds, 63°C for 45 seconds, then 72°C for 75 seconds. After cycling, reactions were finished off with an incubation at 72°C for 5 minutes. The resulting fragment was purified by electrophoresis through a 6% polyacrylamide gel (29:1 cross link) in a buffer of 45 mM Tris-Borate, pH 8.3, 1.4 mM EDTA, visualized by ethidium bromide staining or autoradiography, excised from the gel, eluted by passive diffusion, and concentrated by ethanol precipitation.

Cleavage of the substrate DNA occurred in the presence of the pilot oligonucleotide 19-12 at 50°C (Fig. 12B, lanes 1 and 7) but not at 75°C (lanes 4 and 10). In the presence of oligonucleotide 30-12 cleavage was observed at both temperatures. Cleavage did not occur in the absence of added oligonucleotides (lanes 3, 6 and 12) or at about 80°C even though at 50°C adventitious structures in the substrate allowed primer-independent cleavage in the absence of KCl (Fig. 12B, lane 9). A non-specific oligonucleotide with no complementarity to the substrate DNA did not direct cleavage at 50°C, either in the absence or presence of 50 mM KCl (lanes 13 and 14). Thus, the specificity of the cleavage reactions can be controlled by the extent of complementarity to the substrate and by the conditions of incubation.

#### D. Cleavage Of RNA

An shortened RNA version of the sequence used in the transcleavage experiments discussed above was tested for its ability to serve as a substrate in the reaction. The RNA is cleaved at the expected place, in a reaction that is dependent upon the presence of the pilot oligonucleotide. The RNA substrate, made by T7 RNA polymerase in the presence of [ $\alpha$ - $^{32}$ P]UTP, corresponds to a truncated version of the DNA substrate used in Figure 12B. Reaction conditions were similar to those in used for the DNA substrates described above, with 50 mM KCl; incubation was for 40 minutes at 55°C. The pilot oligonucleotide used is termed 30-0 (SEQ ID NO:20) and is shown in Fig. 13A.

The results of the cleavage reaction is shown in Figure 13B. The reaction was run either in the presence or absence of DNAP*Taq* or pilot oligonucleotide as indicated in Figure 13B.

Strikingly, in the case of RNA cleavage, a 3' arm is not required for the pilot oligonucleotide. It is very unlikely that this cleavage is due to previously described RNaseH, which would be expected to cut the RNA in several places along the 30 base-pair long RNA-DNA duplex. The 5' nuclease of DNAP*Taq* is a structure-specific RNaseH that cleaves the RNA at a single site near the 5' end of the heteroduplexed region.

It is surprising that an oligonucleotide lacking a 3' arm is able to act as a pilot in directing efficient cleavage of an RNA target because such oligonucleotides are unable to direct efficient cleavage of DNA targets using native DNAPs. However, some 5' nucleases of the present invention (for example, clones E, F and G of Figure 4) can cleave DNA in the absence of a 3' arm. In other words, a non-extendable cleavage structure is not required for specific cleavage with some 5' nucleases of the present invention derived from thermostable DNA polymerases.

We tested whether cleavage of an RNA template by DNAP*Taq* in the presence of a fully complementary primer could help explain why DNAP*Taq* is unable to extend a DNA oligonucleotide on an RNA template, in a reaction resembling that of reverse transcriptase. Another thermophilic DNAP, DNAP*Tth*, is able to use RNA as a template, but only in the presence of Mn<sup>++</sup>, so we predicted that this enzyme would not cleave RNA in the presence of this cation.



Accordingly, we incubated an RNA molecule with an appropriate pilot oligonucleotide in the presence of DNAP $Taq$  or DNAP $Tth$ , in buffer containing either  $Mg^{++}$  or  $Mn^{++}$ . As expected, both enzymes cleaved the RNA in the presence of  $Mg^{++}$ . However, DNAP $Taq$ , but not DNAP $Tth$ , degraded the RNA in the presence of  $Mn^{++}$ . We conclude that the 5' nuclease activities of many DNAPs may contribute to their inability to use RNA as templates.

## EXAMPLE 2

### Generation Of 5' Nucleases From Thermostable DNA Polymerases

Thermostable DNA polymerases were generated which have reduced synthetic activity, an activity that is an undesirable side-reaction during DNA cleavage in the detection assay of the invention, yet have maintained thermostable nuclease activity. The result is a thermostable polymerase which cleaves nucleic acids DNA with extreme specificity.

Type A DNA polymerases from eubacteria of the genus *Thermus* share extensive protein sequence identity (90% in the polymerization domain, using the Lipman-Pearson method in the DNA analysis software from DNASTar, WI) and behave similarly in both polymerization and nuclease assays. Therefore, we have used the genes for the DNA polymerase of *Thermus aquaticus* (DNAP $Taq$ ) and *Thermus flavus* (DNAP $Tfl$ ) as representatives of this class. Polymerase genes from other eubacterial organisms, such as *Thermus thermophilus*, *Thermus sp.*, *Thermotoga maritima*, *Thermosipho africanus* and *Bacillus stearothermophilus* are equally suitable. The DNA polymerases from these thermophilic organisms are capable of surviving and performing at elevated temperatures, and can thus be used in reactions in which temperature is used as a selection against non-specific hybridization of nucleic acid strands.

The restriction sites used for deletion mutagenesis, described below, were chosen for convenience. Different sites situated with similar convenience are available in the *Thermus thermophilus* gene and can be used to make similar constructs with other Type A polymerase genes from related organisms.

## A. Creation Of 5' Nuclease Constructs

### 1. Modified DNAP $Taq$ Genes

The first step was to place a modified gene for the  $Taq$  DNA polymerase on a plasmid under control of an inducible promoter. The modified  $Taq$  polymerase gene was isolated as follows: The  $Taq$  DNA polymerase gene was amplified by polymerase chain reaction from genomic DNA from *Thermus aquaticus*, strain YT-1 (Lawyer *et al.*, *supra*), using as primers the oligonucleotides described in SEQ ID NOS:13-14. The resulting fragment of DNA has a recognition sequence for the restriction endonuclease EcoRI at the 5' end of the coding sequence and a BglII sequence at the 3' end. Cleavage with BglII leaves a 5' overhang or "sticky end" that is compatible with the end generated by BamHI. The PCR-amplified DNA was digested with EcoRI and BamHI. The 2512 bp fragment containing the coding region for the polymerase gene was gel purified and then ligated into a plasmid which contains an inducible promoter.

In one embodiment of the invention, the pTTQ18 vector, which contains the hybrid *trp-lac* (*tac*) promoter, was used [M.J.R. Stark, *Gene* 5:255 (1987)] and shown in Fig. 14. The *tac* promoter is under the control of the *E. coli lac* repressor. Repression allows the synthesis of the gene product to be suppressed until the desired level of bacterial growth has been achieved, at which point repression is removed by addition of a specific inducer, isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG). Such a system allows the expression of foreign proteins that may slow or prevent growth of transformants.

Bacterial promoters, such as *tac*, may not be adequately suppressed when they are present on a multiple copy plasmid. If a highly toxic protein is placed under control of such a promoter, the small amount of expression leaking through can be harmful to the bacteria. In another embodiment of the invention, another option for repressing synthesis of a cloned gene product was used. The non-bacterial promoter, from bacteriophage T7, found in the plasmid vector series pET-3 was used to express the cloned mutant  $Taq$  polymerase genes [Fig. 15; Studier and Moffatt, *J. Mol. Biol.* 189:113 (1986)]. This promoter initiates transcription only by T7 RNA polymerase. In a suitable strain, such as BL21(DE3)pLYS, the gene for this RNA polymerase is carried on the bacterial genome under control of

the *lac* operator. This arrangement has the advantage that expression of the multiple copy gene (on the plasmid) is completely dependent on the expression of T7 RNA polymerase, which is easily suppressed because it is present in a single copy.

5           For ligation into the pTTQ18 vector (Fig. 14), the PCR product DNA containing the *Taq* polymerase coding region (*mutTaq*, clone 4B, SEQ ID NO:21) was digested with EcoRI and BglII and this fragment was ligated under standard "sticky end" conditions [Sambrook *et al. Molecular Cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1989)] into the EcoRI and BamHI sites of  
10           the plasmid vector pTTQ18. Expression of this construct yields a translational fusion product in which the first two residues of the native protein (Met-Arg) are replaced by three from the vector (Met-Asn-Ser), but the remainder of the natural protein would not change. The construct was transformed into the JM109 strain of *E. coli* and the transformants were plated under incompletely repressing conditions  
15           that do not permit growth of bacteria expressing the native protein. These plating conditions allow the isolation of genes containing pre-existing mutations, such as those that result from the infidelity of *Taq* polymerase during the amplification process.

          Using this amplification/selection protocol, we isolated a clone (depicted in  
20           Fig. 4B) containing a mutated *Taq* polymerase gene (*mutTaq*, clone 4B). The mutant was first detected by its phenotype, in which temperature-stable 5' nuclease activity in a crude cell extract was normal, but polymerization activity was almost absent (approximately less than 1% of wild type *Taq* polymerase activity).

          DNA sequence analysis of the recombinant gene showed that it had changes  
25           in the polymerase domain resulting in two amino acid substitutions: an A to G change at nucleotide position 1394 causes a Glu to Gly change at amino acid position 465 (numbered according to the natural nucleic and amino acid sequences, SEQ ID NOS:1 and 4) and another A to G change at nucleotide position 2260  
          causes a Gln to Arg change at amino acid position 754. Because the Gln to Gly  
30           mutation is at a nonconserved position and because the Glu to Arg mutation alters an amino acid that is conserved in virtually all of the known Type A polymerases, this latter mutation is most likely the one responsible for curtailing the synthesis

activity of this protein. The nucleotide sequence for the Fig. 4B construct is given in SEQ ID NO:21.

Subsequent derivatives of DNAP*Taq* constructs were made from the mut*Taq* gene, thus, they all bear these amino acid substitutions in addition to their other alterations, unless these particular regions were deleted. These mutated sites are indicated by black boxes at these locations in the diagrams in Fig. 4. All constructs except the genes shown in Figures 4E, F and G were made in the pTTQ18 vector.

The cloning vector used for the genes in Figs. 4E and F was from the commercially available pET-3 series, described above. Though this vector series has only a BamHI site for cloning downstream of the T7 promoter, the series contains variants that allow cloning into any of the three reading frames. For cloning of the PCR product described above, the variant called pET-3c was used (Fig 15). The vector was digested with BamHI, dephosphorylated with calf intestinal phosphatase, and the sticky ends were filled in using the Klenow fragment of DNAPEc1 and dNTPs. The gene for the mutant *Taq* DNAP shown in Fig. 4B (mut*Taq*, clone 4B) was released from pTTQ18 by digestion with EcoRI and SalI, and the "sticky ends" were filled in as was done with the vector. The fragment was ligated to the vector under standard blunt-end conditions (Sambrook *et al.*, *Molecular Cloning, supra*), the construct was transformed into the BL21(DE3)pLYS strain of *E. coli*, and isolates were screened to identify those that were ligated with the gene in the proper orientation relative to the promoter. This construction yields another translational fusion product, in which the first two amino acids of DNAP*Taq* (Met-Arg) are replaced by 13 from the vector plus two from the PCR primer (Met-Ala-Ser-Met-Thr-Gly-Gly-Gln-Gln-Met-Gly-Arg-Ile-Asn-Ser) (SEQ ID NO:29).

Our goal was to generate enzymes that lacked the ability to synthesize DNA, but retained the ability to cleave nucleic acids with a 5' nuclease activity. The act of primed, templated synthesis of DNA is actually a coordinated series of events, so it is possible to disable DNA synthesis by disrupting one event while not affecting the others. These steps include, but are not limited to, primer recognition and binding, dNTP binding and catalysis of the inter-nucleotide phosphodiester

bond. Some of the amino acids in the polymerization domain of DNAPEcI have been linked to these functions, but the precise mechanisms are as yet poorly defined.

5 One way of destroying the polymerizing ability of a DNA polymerase is to delete all or part of the gene segment that encodes that domain for the protein, or to otherwise render the gene incapable of making a complete polymerization domain. Individual mutant enzymes may differ from each other in stability and solubility both inside and outside cells. For instance, in contrast to the 5' nuclease domain of DNAPEcI, which can be released in an active form from the  
10 polymerization domain by gentle proteolysis [Setlow and Kornberg, *J. Biol. Chem.* 247:232 (1972)], the *Thermus* nuclease domain, when treated similarly, becomes less soluble and the cleavage activity is often lost.

Using the mutant gene shown in Fig. 4B as starting material, several deletion constructs were created. All cloning technologies were standard  
15 (Sambrook *et al.*, *supra*) and are summarized briefly, as follows:

Fig. 4C: The mut*Taq* construct was digested with PstI, which cuts once within the polymerase coding region, as indicated, and cuts immediately downstream of the gene in the multiple cloning site of the vector. After release of the fragment between these two sites, the vector was re-ligated, creating an 894-  
20 nucleotide deletion, and bringing into frame a stop codon 40 nucleotides downstream of the junction. The nucleotide sequence of this 5' nuclease (clone 4C) is given in SEQ ID NO:9.

Fig. 4D: The mut*Taq* construct was digested with NheI, which cuts once in the gene at position 2047. The resulting four-nucleotide 5' overhanging ends were  
25 filled in, as described above, and the blunt ends were re-ligated. The resulting four-nucleotide insertion changes the reading frame and causes termination of translation ten amino acids downstream of the mutation. The nucleotide sequence of this 5' nuclease (clone 4D) is given in SEQ ID NO:10.

Fig. 4E: The entire mut*Taq* gene was cut from pTTQ18 using EcoRI and  
30 SalI and cloned into pET-3c, as described above. This clone was digested with BstXI and XcmI, at unique sites that are situated as shown in Fig. 4E. The DNA was treated with the Klenow fragment of DNAPEcI and dNTPs, which resulted in

the 3' overhangs of both sites being trimmed to blunt ends. These blunt ends were ligated together, resulting in an out-of-frame deletion of 1540 nucleotides. An in-frame termination codon occurs 18 triplets past the junction site. The nucleotide sequence of this 5' nuclease (clone 4E) is given in SEQ ID NO:11, with the appropriate leader sequence given in SEQ ID NO:30. It is also referred to as Cleavase™ BX.

Fig. 4F: The entire *mutTaq* gene was cut from pTTQ18 using EcoRI and Sall and cloned into pET-3c, as described above. This clone was digested with BstXI and amHI, at unique sites that are situated as shown in the diagram. The DNA was treated with the Klenow fragment of DNAPEc1 and dNTPs, which resulted in the 3' overhang of the BstX I site being trimmed to a blunt end, while the 5' overhang of the Bam HI site was filled in to make a blunt end. These ends were ligated together, resulting in an in-frame deletion of 903 nucleotides. The nucleotide sequence of the 5' nuclease (clone 4F) is given in SEQ ID NO:12. It is also referred to as Cleavase™ BB.

Fig.4G: This polymerase is a variant of that shown in Figure 4E. It was cloned in the plasmid vector pET-21 (Novagen). The non-bacterial promoter from bacteriophage T7, found in this vector, initiates transcription only by T7 RNA polymerase. See Studier and Moffatt, *supra*. In a suitable strain, such as (DES)pLYS, the gene for this RNA polymerase is carried on the bacterial genome under control of the *lac* operator. This arrangement has the advantage that expression of the multiple copy gene (on the plasmid) is completely dependent on the expression of T7 RNA polymerase, which is easily suppressed because it is present in a single copy. Because the expression of these mutant genes is under this tightly controlled promoter, potential problems of toxicity of the expressed proteins to the host cells are less of a concern.

The pET-21 vector also features a "His\*Tag", a stretch of six consecutive histidine residues that are added on the carboxy terminus of the expressed proteins. The resulting proteins can then be purified in a single step by metal chelation chromatography, using a commercially available (Novagen) column resin with immobilized Ni<sup>++</sup> ions. The 2.5 ml columns are reusable, and can bind up to 20



mg of the target protein under native or denaturing (guanidine\*HCl or urea) conditions.

5 *E. coli* (DES)pLYS cells are transformed with the constructs described above using standard transformation techniques, and used to inoculate a standard growth medium (*e.g.*, Luria-Bertani broth). Production of T7 RNA polymerase is induced during log phase growth by addition of IPTG and incubated for a further 12 to 17 hours. Aliquots of culture are removed both before and after induction and the proteins are examined by SDS-PAGE. Staining with Coomassie Blue allows visualization of the foreign proteins if they account for about 3-5% of the cellular protein and do not co-migrate with any of the major protein bands. 10 Proteins that co-migrate with major host protein must be expressed as more than 10% of the total protein to be seen at this stage of analysis.

Some mutant proteins are sequestered by the cells into inclusion bodies. These are granules that form in the cytoplasm when bacteria are made to express 15 high levels of a foreign protein, and they can be purified from a crude lysate, and analyzed by SDS-PAGE to determine their protein content. If the cloned protein is found in the inclusion bodies, it must be released to assay the cleavage and polymerase activities. Different methods of solubilization may be appropriate for different proteins, and a variety of methods are known. *See e.g.* Builder & Ogez, 20 U.S. Patent No. 4,511,502 (1985); Olson, U.S. Patent No. 4,518,526 (1985); Olson & Pai, U.S. Patent No. 4,511,503 (1985); Jones *et al.*, U.S. Patent No. 4,512,922 (1985), all of which are hereby incorporated by reference.

The solubilized protein is then purified on the Ni<sup>++</sup> column as described above, following the manufacturers instructions (Novagen). The washed proteins 25 are eluted from the column by a combination of imidazole competitor (1 M) and high salt (0.5 M NaCl), and dialyzed to exchange the buffer and to allow denature proteins to refold. Typical recoveries result in approximately 20 µg of specific protein per ml of starting culture. The DNAP mutant is referred to as Cleavase™ BN and the sequence is given in SEQ ID NO:31.

## 2. Modified DNAPTfl Gene

The DNA polymerase gene of *Thermus flavus* was isolated from the "T. *flavus*" AT-62 strain obtained from the American Type Tissue Collection (ATCC 33923). This strain has a different restriction map than does the *T. flavus* strain used to generate the sequence published by Akhmetzjanov and Vakhitov, *supra*. The published sequence is listed as SEQ ID NO:2. No sequence data has been published for the DNA polymerase gene from the AT-62 strain of *T. flavus*.

Genomic DNA from *T. flavus* was amplified using the same primers used to amplify the *T. aquaticus* DNA polymerase gene (SEQ ID NOS:13-14). The approximately 2500 base pair PCR fragment was digested with EcoRI and BamHI. The over-hanging ends were made blunt with the Klenow fragment of DNAPEc1 and dNTPs. The resulting approximately 1800 base pair fragment containing the coding region for the N-terminus was ligated into pET-3c, as described above. This construct, clone 5B, is depicted in Fig. 5B. The wild type *T. flavus* DNA polymerase gene is depicted in Fig. 5A. The 5B clone has the same leader amino acids as do the DNAPTaq clones 4E and F which were cloned into pET-3c; it is not known precisely where translation termination occurs, but the vector has a strong transcription termination signal immediately downstream of the cloning site.

### B. Growth And Induction Of Transformed Cells

Bacterial cells were transformed with the constructs described above using standard transformation techniques and used to inoculate 2 mls of a standard growth medium (*e.g.*, Luria-Bertani broth). The resulting cultures were incubated as appropriate for the particular strain used, and induced if required for a particular expression system. For all of the constructs depicted in Figs. 4 and 5, the cultures were grown to an optical density (at 600nm wavelength) of 0.5 OD.

To induce expression of the cloned genes, the cultures were brought to a final concentration of 0.4 mM IPTG and the incubations were continued for 12 to 17 hours. 50  $\mu$ l aliquots of each culture were removed both before and after induction and were combined with 20  $\mu$ l of a standard gel loading buffer for sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Subsequent staining with Coomassie Blue (Sambrook *et al.*, *supra*) allows

visualization of the foreign proteins if they account for about 3-5% of the cellular protein and do not co-migrate with any of the major *E. coli* protein bands.

Proteins that do co-migrate with a major host protein must be expressed as more than 10% of the total protein to be seen at this stage of analysis.

### 5 C. Heat Lysis And Fractionation

Expressed thermostable proteins, *i.e.*, the 5' nucleases, were isolated by heating crude bacterial cell extracts to cause denaturation and precipitation of the less stable *E. coli* proteins. The precipitated *E. coli* proteins were then, along with other cell debris, removed by centrifugation. 1.7 mls of the culture were pelleted  
10 by microcentrifugation at 12,000 to 14,000 rpm for 30 to 60 seconds. After removal of the supernatant, the cells were resuspended in 400  $\mu$ l of buffer A (50 mM Tris-HCl, pH 7.9, 50 mM dextrose, 1 mM EDTA), re-centrifuged, then resuspended in 80  $\mu$ l of buffer A with 4mg/ml lysozyme. The cells were incubated at room temperature for 15 minutes, then combined with 80  $\mu$ l of buffer B (10 mM  
15 Tris-HCl, pH 7.9, 50 mM KCl, 1 mM EDTA, 1 mM PMSF, 0.5% Tween-20, 0.5% Nonidet-P40).

This mixture was incubated at 75°C for 1 hour to denature and precipitate the host proteins. This cell extract was centrifuged at 14,000 rpm for 15 minutes at 4°C, and the supernatant was transferred to a fresh tube. An aliquot of 0.5 to 1  $\mu$ l  
20 of this supernatant was used directly in each test reaction, and the protein content of the extract was determined by subjecting 7  $\mu$ l to electrophoretic analysis, as above. The native recombinant *Taq* DNA polymerase [Englke, Anal. Biochem 191:396 (1990)], and the double point mutation protein shown in Fig. 4B are both soluble and active at this point.

25 The foreign protein may not be detected after the heat treatments due to sequestration of the foreign protein by the cells into inclusion bodies. These are granules that form in the cytoplasm when bacteria are made to express high levels of a foreign protein, and they can be purified from a crude lysate, and analyzed SDS PAGE to determine their protein content. Many methods have been described  
30 in the literature, and one approach is described below.

#### D. Isolation And Solubilization Of Inclusion Bodies

A small culture was grown and induced as described above. A 1.7 ml aliquot was pelleted by brief centrifugation, and the bacterial cells were resuspended in 100  $\mu$ l of Lysis buffer (50 mM Tris-HCl, pH 8.0, 1 mM EDTA, 100 mM NaCl). 2.5  $\mu$ l of 20 mM PMSF were added for a final concentration of 0.5 mM, and lysozyme was added to a concentration of 1.0 mg/ml. The cells were incubated at room temperature for 20 minutes, deoxycholic acid was added to 1mg/ml (1  $\mu$ l of 100 mg/ml solution), and the mixture was further incubated at 37°C for about 15 minutes or until viscous. DNase I was added to 10  $\mu$ g/ml and the mixture was incubated at room temperature for about 30 minutes or until it was no longer viscous.

From this mixture the inclusion bodies were collected by centrifugation at 14,000 rpm for 15 minutes at 4°C, and the supernatant was discarded. The pellet was resuspended in 100  $\mu$ l of lysis buffer with 10mM EDTA (pH 8.0) and 0.5% Triton X-100. After 5 minutes at room temperature, the inclusion bodies were pelleted as before, and the supernatant was saved for later analysis. The inclusion bodies were resuspended in 50  $\mu$ l of distilled water, and 5  $\mu$ l was combined with SDS gel loading buffer (which dissolves the inclusion bodies) and analyzed electrophoretically, along with an aliquot of the supernatant.

If the cloned protein is found in the inclusion bodies, it may be released to assay the cleavage and polymerase activities and the method of solubilization must be compatible with the particular activity. Different methods of solubilization may be appropriate for different proteins, and a variety of methods are discussed in *Molecular Cloning* (Sambrook *et al.*, *supra*). The following is an adaptation we have used for several of our isolates.

20  $\mu$ l of the inclusion body-water suspension were pelleted by centrifugation at 14,000 rpm for 4 minutes at room temperature, and the supernatant was discarded. To further wash the inclusion bodies, the pellet was resuspended in 20 $\mu$ l of lysis buffer with 2M urea, and incubated at room temperature for one hour. The washed inclusion bodies were then resuspended in 2  $\mu$ l of lysis buffer with 8M urea; the solution clarified visibly as the inclusion bodies dissolved. Undissolved

debris was removed by centrifugation at 14,000 rpm for 4 minutes at room temperature, and the extract supernatant was transferred to a fresh tube.

To reduce the urea concentration, the extract was diluted into  $\text{KH}_2\text{PO}_4$ . A fresh tube was prepared containing 180  $\mu\text{l}$  of 50 mM  $\text{KH}_2\text{PO}_4$ , pH 9.5, 1 mM EDTA and 50 mM NaCl. A 2  $\mu\text{l}$  aliquot of the extract was added and vortexed briefly to mix. This step was repeated until all of the extract had been added for a total of 10 additions. The mixture was allowed to sit at room temperature for 15 minutes, during which time some precipitate often forms. Precipitates were removed by centrifugation at 14,000 rpm, for 15 minutes at room temperature, and the supernatant was transferred to a fresh tube. To the 200  $\mu\text{l}$  of protein in the  $\text{KH}_2\text{PO}_4$  solution, 140-200  $\mu\text{l}$  of saturated  $(\text{NH}_4)_2\text{SO}_4$  were added, so that the resulting mixture was about 41% to 50% saturated  $(\text{NH}_4)_2\text{SO}_4$ . The mixture was chilled on ice for 30 minutes to allow the protein to precipitate, and the protein was then collected by centrifugation at 14,000 rpm, for 4 minutes at room temperature. The supernatant was discarded, and the pellet was dissolved in 20  $\mu\text{l}$  Buffer C (20 mM HEPES, pH 7.9, 1 mM EDTA, 0.5% PMSF, 25 mM KCl and 0.5 % each of Tween-20 and Nonidet P 40). The protein solution was centrifuged again for 4 minutes to pellet insoluble materials, and the supernatant was removed to a fresh tube. The protein contents of extracts prepared in this manner were visualized by resolving 1-4  $\mu\text{l}$  by SDS-PAGE; 0.5 to 1  $\mu\text{l}$  of extract was tested in the cleavage and polymerization assays as described.

#### **E. Protein Analysis For Presence Of Nuclease And Synthetic Activity**

The 5' nucleases described above and shown in Figs. 4 and 5 were analyzed by the following methods.

##### **1. Structure Specific Nuclease Assay**

A candidate modified polymerase is tested for 5' nuclease activity by examining its ability to catalyze structure-specific cleavages. By the term "cleavage structure" as used herein, is meant a nucleic acid structure which is a substrate for cleavage by the 5' nuclease activity of a DNAP.

5 The polymerase is exposed to test complexes that have the structures shown in Fig. 16. Testing for 5' nuclease activity involves three reactions: 1) a primer-directed cleavage (Fig. 16B) is performed because it is relatively insensitive to variations in the salt concentration of the reaction and can, therefore, be performed in whatever solute conditions the modified enzyme requires for activity; this is generally the same conditions preferred by unmodified polymerases; 2) a similar primer-directed cleavage is performed in a buffer which permits primer-independent cleavage, *i.e.*, a low salt buffer, to demonstrate that the enzyme is viable under these conditions; and 3) a primer-independent cleavage (Fig. 16A) is performed in the same low salt buffer.

10 The bifurcated duplex is formed between a substrate strand and a template strand as shown in Fig. 16. By the term "substrate strand" as used herein, is meant that strand of nucleic acid in which the cleavage mediated by the 5' nuclease activity occurs. The substrate strand is always depicted as the top strand in the bifurcated complex which serves as a substrate for 5' nuclease cleavage (Fig. 16). By the term "template strand" as used herein, is meant the strand of nucleic acid which is at least partially complementary to the substrate strand and which anneals to the substrate strand to form the cleavage structure. The template strand is always depicted as the bottom strand of the bifurcated cleavage structure (Fig. 16). If a primer (a short oligonucleotide of 19 to 30 nucleotides in length) is added to the complex, as when primer-dependent cleavage is to be tested, it is designed to anneal to the 3' arm of the template strand (Fig. 16B). Such a primer would be extended along the template strand if the polymerase used in the reaction has synthetic activity.

25 The cleavage structure may be made as a single hairpin molecule, with the 3' end of the target and the 5' end of the pilot joined as a loop as shown in Fig. 16E. A primer oligonucleotide complementary to the 3' arm is also required for these tests so that the enzyme's sensitivity to the presence of a primer may be tested.

30 Nucleic acids to be used to form test cleavage structures can be chemically synthesized, or can be generated by standard recombinant DNA techniques. By the latter method, the hairpin portion of the molecule can be created by inserting into a



cloning vector duplicate copies of a short DNA segment, adjacent to each other but in opposing orientation. The double-stranded fragment encompassing this inverted repeat, and including enough flanking sequence to give short (about 20 nucleotides) unpaired 5' and 3' arms, can then be released from the vector by restriction enzyme digestion, or by PCR performed with an enzyme lacking a 5' exonuclease (e.g., the Stoffel fragment of Amplitaq™ DNA polymerase, Vent™ DNA polymerase).

The test DNA can be labeled on either end, or internally, with either a radioisotope, or with a non-isotopic tag. Whether the hairpin DNA is a synthetic single strand or a cloned double strand, the DNA is heated prior to use to melt all duplexes. When cooled on ice, the structure depicted in Fig. 16E is formed, and is stable for sufficient time to perform these assays.

To test for primer-directed cleavage (Reaction 1), a detectable quantity of the test molecule (typically 1-100 fmol of <sup>32</sup>P-labeled hairpin molecule) and a 10 to 100-fold molar excess of primer are placed in a buffer known to be compatible with the test enzyme. For Reaction 2, where primer-directed cleavage is performed under condition which allow primer-independent cleavage, the same quantities of molecules are placed in a solution that is the same as the buffer used in Reaction 1 regarding pH, enzyme stabilizers (e.g., bovine serum albumin, nonionic detergents, gelatin) and reducing agents (e.g., dithiothreitol, 2-mercaptoethanol) but that replaces any monovalent cation salt with 20 mM KCl; 20 mM KCl is the demonstrated optimum for primer-independent cleavage. Buffers for enzymes, such as DNAPEc1, that usually operate in the absence of salt are not supplemented to achieve this concentration. To test for primer-independent cleavage (Reaction 3) the same quantity of the test molecule, but no primer, are combined under the same buffer conditions used for Reaction 2.

All three test reactions are then exposed to enough of the enzyme that the molar ratio of enzyme to test complex is approximately 1:1. The reactions are incubated at a range of temperatures up to, but not exceeding, the temperature allowed by either the enzyme stability or the complex stability, whichever is lower, up to 80°C for enzymes from thermophiles, for a time sufficient to allow cleavage (10 to 60 minutes). The products of Reactions 1, 2 and 3 are resolved by

denaturing polyacrylamide gel electrophoresis, and visualized by autoradiography or by a comparable method appropriate to the labeling system used. Additional labeling systems include chemiluminescence detection, silver or other stains, blotting and probing and the like. The presence of cleavage products is indicated by the presence of molecules which migrate at a lower molecular weight than does the uncleaved test structure. These cleavage products indicate that the candidate polymerase has structure-specific 5' nuclease activity.

To determine whether a modified DNA polymerase has substantially the same 5' nuclease activity as that of the native DNA polymerase, the results of the above-described tests are compared with the results obtained from these tests performed with the native DNA polymerase. By "substantially the same 5' nuclease activity" we mean that the modified polymerase and the native polymerase will both cleave test molecules in the same manner. It is not necessary that the modified polymerase cleave at the same rate as the native DNA polymerase.

Some enzymes or enzyme preparations may have other associated or contaminating activities that may be functional under the cleavage conditions described above and that may interfere with 5' nuclease detection. Reaction conditions can be modified in consideration of these other activities, to avoid destruction of the substrate, or other masking of the 5' nuclease cleavage and its products. For example, the DNA polymerase I of *E. coli* (Pol I), in addition to its polymerase and 5' nuclease activities, has a 3' exonuclease that can degrade DNA in a 3' to 5' direction. Consequently, when the molecule in Fig. 16E is exposed to this polymerase under the conditions described above, the 3' exonuclease quickly removes the unpaired 3' arm, destroying the bifurcated structure required of a substrate for the 5' exonuclease cleavage and no cleavage is detected. The true ability of Pol I to cleave the structure can be revealed if the 3' exonuclease is inhibited by a change of conditions (*e.g.*, pH), mutation, or by addition of a competitor for the activity. Addition of 500 pmoles of a single-stranded competitor oligonucleotide, unrelated to the Fig. 16E structure, to the cleavage reaction with Pol I effectively inhibits the digestion of the 3' arm of the Fig. 16E structure without interfering with the 5' exonuclease release of the 5' arm. The

concentration of the competitor is not critical, but should be high enough to occupy the 3' exonuclease for the duration of the reaction.

Similar destruction of the test molecule may be caused by contaminants in the candidate polymerase preparation. Several sets of the structure specific nuclease reactions may be performed to determine the purity of the candidate nuclease and to find the window between under and over exposure of the test molecule to the polymerase preparation being investigated.

The above described modified polymerases were tested for 5' nuclease activity as follows: Reaction 1 was performed in a buffer of 10 mM Tris-Cl, pH 8.5 at 20°C, 1.5 mM MgCl<sub>2</sub> and 50 mM KCl and in Reaction 2 the KCl concentration was reduced to 20 mM. In Reactions 1 and 2, 10 fmoles of the test substrate molecule shown in Fig. 16E were combined with 1 pmole of the indicated primer and 0.5 to 1.0 µl of extract containing the modified polymerase (prepared as described above). This mixture was then incubated for 10 minutes at 55°C. For all of the mutant polymerases tested these conditions were sufficient to give complete cleavage. When the molecule shown in Fig. 16E was labeled at the 5' end, the released 5' fragment, 25 nucleotides long, was conveniently resolved on a 20% polyacrylamide gel (19:1 cross-linked) with 7 M urea in a buffer containing 45 mM Tris-borate pH 8.3, 1.4 mM EDTA. Clones 4C-F and 5B exhibited structure-specific cleavage comparable to that of the unmodified DNA polymerase. Additionally, clones 4E, 4F and 4G have the added ability to cleave DNA in the absence of a 3' arm as discussed above. Representative cleavage reactions are shown in Figure 17.

For the reactions shown in Fig. 17, the mutant polymerase clones 4E (*Taq* mutant) and 5B (*Tfl* mutant) were examined for their ability to cleave the hairpin substrate molecule shown in Fig. 16E. The substrate molecule was labeled at the 5' terminus with <sup>32</sup>P. 10 fmoles of heat-denatured, end-labeled substrate DNA and 0.5 units of DNAP*Taq* (lane 1) or 0.5 µl of 4e or 5b extract (Fig. 17, lanes 2-7, extract was prepared as described above) were mixed together in a buffer containing 10 mM Tris-Cl, pH 8.5, 50 mM KCl and 1.5 mM MgCl<sub>2</sub>. The final reaction volume was 10 µl. Reactions shown in lanes 4 and 7 contain in addition 50 µM of each dNTP. Reactions shown in lanes 3, 4, 6 and 7 contain 0.2 µM of

the primer oligonucleotide (complementary to the 3' arm of the substrate and shown in Fig. 16E). Reactions were incubated at 55° C for 4 minutes. Reactions were stopped by the addition of 8 µl of 95% formamide containing 20 mM EDTA and 0.05% marker dyes per 10 µl reaction volume. Samples were then applied to 12% denaturing acrylamide gels. Following electrophoresis, the gels were audioradiographed. Fig. 17 shows that clones 4E and 5B exhibit cleavage activity similar to that of the native DNAP $Taq$ . Note that some cleavage occurs in these reactions in the absence of the primer. When long hairpin structure, such as the one used here (Fig. 16E), are used in cleavage reactions performed in buffers containing 50 mM KCl a low level of primer-independent cleavage is seen. Higher concentrations of KCl suppress, but do not eliminate, this primer-independent cleavage under these conditions.

## 2. Assay For Synthetic Activity

The ability of the modified enzyme or proteolytic fragments is assayed by adding the modified enzyme to an assay system in which a primer is annealed to a template and DNA synthesis is catalyzed by the added enzyme. Many standard laboratory techniques employ such an assay. For example, nick translation and enzymatic sequencing involve extension of a primer along a DNA template by a polymerase molecule.

In a preferred assay for determining the synthetic activity of a modified enzyme an oligonucleotide primer is annealed to a single-stranded DNA template, *e.g.*, bacteriophage M13 DNA, and the primer/template duplex is incubated in the presence of the modified polymerase in question, deoxynucleoside triphosphates (dNTPs) and the buffer and salts known to be appropriate for the unmodified or native enzyme. Detection of either primer extension (by denaturing gel electrophoresis) or dNTP incorporation (by acid precipitation or chromatography) is indicative of an active polymerase. A label, either isotopic or non-isotopic, is preferably included on either the primer or as a dNTP to facilitate detection of polymerization products. Synthetic activity is quantified as the amount of free nucleotide incorporated into the growing DNA chain and is expressed as amount incorporated per unit of time under specific reaction conditions.

Representative results of an assay for synthetic activity is shown in Fig. 18. The synthetic activity of the mutant DNAP*Taq* clones 4B-F was tested as follows: A master mixture of the following buffer was made: 1.2X PCR buffer (1X PCR buffer contains 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 10 mM Tris-Cl, pH 8.5 and 0.05% each Tween 20 and Nonidet P40), 50 μM each of dGTP, dATP and dTTP, 5 μM dCTP and 0.125 μM α-<sup>32</sup>P-dCTP at 600 Ci/mmol. Before adjusting this mixture to its final volume, it was divided into two equal aliquots. One received distilled water up to a volume of 50 μl to give the concentrations above. The other received 5 μg of single-stranded M13mp18 DNA (approximately 2.5 pmol or 0.05 μM final concentration) and 250 pmol of M13 sequencing primer (5 μM final concentration) and distilled water to a final volume of 50 μl. Each cocktail was warmed to 75°C for 5 minutes and then cooled to room temperature. This allowed the primers to anneal to the DNA in the DNA-containing mixtures.

For each assay, 4 μl of the cocktail with the DNA was combined with 1 μl of the mutant polymerase, prepared as described, or 1 unit of DNAP*Taq* (Perkin Elmer) in 1 μl of dH<sub>2</sub>O. A "no DNA" control was done in the presence of the DNAP*Taq* (Fig. 18, lane 1), and a "no enzyme" control was done using water in place of the enzyme (lane 2). Each reaction was mixed, then incubated at room temperature (approx. 22°C) for 5 minutes, then at 55°C for 2 minutes, then at 72°C for 2 minutes. This step incubation was done to detect polymerization in any mutants that might have optimal temperatures lower than 72°C. After the final incubation, the tubes were spun briefly to collect any condensation and were placed on ice. One μl of each reaction was spotted at an origin 1.5 cm from the bottom edge of a polyethyleneimine (PEI) cellulose thin layer chromatography plate and allowed to dry. The chromatography plate was run in 0.75 M NaH<sub>2</sub>PO<sub>4</sub>, pH 3.5, until the buffer front had run approximately 9 cm from the origin. The plate was dried, wrapped in plastic wrap, marked with luminescent ink, and exposed to X-ray film. Incorporation was detected as counts that stuck where originally spotted, while the unincorporated nucleotides were carried by the salt solution from the origin.

Comparison of the locations of the counts with the two control lanes confirmed the lack of polymerization activity in the mutant preparations. Among



the modified DNAP*Taq* clones, only clone 4B retains any residual synthetic activity as shown in Fig. 18.

### EXAMPLE 3

#### 5' Nucleases Derived From Thermostable DNA Polymerases Can Cleave Short Hairpin Structures With Specificity

The ability of the 5' nucleases to cleave hairpin structures to generate a cleaved hairpin structure suitable as a detection molecule was examined. The structure and sequence of the hairpin test molecule is shown in Fig. 19A (SEQ ID NO:15). The oligonucleotide (labeled "primer" in Fig. 19A, SEQ ID NO:22) is shown annealed to its complementary sequence on the 3' arm of the hairpin test molecule. The hairpin test molecule was single-end labeled with  $^{32}\text{P}$  using a labeled T7 promoter primer in a polymerase chain reaction. The label is present on the 5' arm of the hairpin test molecule and is represented by the star in Fig. 19A.

The cleavage reaction was performed by adding 10 fmoles of heat-denatured, end-labeled hairpin test molecule, 0.2  $\mu\text{M}$  of the primer oligonucleotide (complementary to the 3' arm of the hairpin), 50  $\mu\text{M}$  of each dNTP and 0.5 units of DNAP*Taq* (Perkin Elmer) or 0.5  $\mu\text{l}$  of extract containing a 5' nuclease (prepared as described above) in a total volume of 10  $\mu\text{l}$  in a buffer containing 10 mM Tris-Cl, pH 8.5, 50 mM KCl and 1.5 mM  $\text{MgCl}_2$ . Reactions shown in lanes 3, 5 and 7 were run in the absence of dNTPs.

Reactions were incubated at 55° C for 4 minutes. Reactions were stopped at 55° C by the addition of 8  $\mu\text{l}$  of 95% formamide with 20 mM EDTA and 0.05% marker dyes per 10  $\mu\text{l}$  reaction volume. Samples were not heated before loading onto denaturing polyacrylamide gels (10% polyacrylamide, 19:1 crosslinking, 7 M urea, 89 mM Tris-borate, pH 8.3, 2.8 mM EDTA). The samples were not heated to allow for the resolution of single-stranded and re-duplexed uncleaved hairpin molecules.

Fig. 19B shows that altered polymerases lacking any detectable synthetic activity cleave a hairpin structure when an oligonucleotide is annealed to the single-stranded 3' arm of the hairpin to yield a single species of cleaved product (Fig.



19B, lanes 3 and 4). 5' nucleases, such as clone 4D, shown in lanes 3 and 4, produce a single cleaved product even in the presence of dNTPs. 5' nucleases which retain a residual amount of synthetic activity (less than 1% of wild type activity) produce multiple cleavage products as the polymerase can extend the oligonucleotide annealed to the 3' arm of the hairpin thereby moving the site of cleavage (clone 4B, lanes 5 and 6). Native DNA*Taq* produces even more species of cleavage products than do mutant polymerases retaining residual synthetic activity and additionally converts the hairpin structure to a double-stranded form in the presence of dNTPs due to the high level of synthetic activity in the native polymerase (Fig. 19B, lane 8).

#### EXAMPLE 4

##### Test Of The Trigger/Detection Assay

To test the ability of an oligonucleotide of the type released in the trigger reaction of the trigger/detection assay to be detected in the detection reaction of the assay, the two hairpin structures shown in Fig. 20A were synthesized using standard techniques. The two hairpins are termed the A-hairpin (SEQ ID NO:23) and the T-hairpin (SEQ ID NO:24). The predicted sites of cleavage in the presence of the appropriate annealed primers are indicated by the arrows. The A- and T-hairpins were designed to prevent intra-strand mis-folding by omitting most of the T residues in the A-hairpin and omitting most of the A residues in the T-hairpin. To avoid mis-priming and slippage, the hairpins were designed with local variations in the sequence motifs (*e.g.*, spacing T residues one or two nucleotides apart or in pairs). The A- and T-hairpins can be annealed together to form a duplex which has appropriate ends for directional cloning in pUC-type vectors; restriction sites are located in the loop regions of the duplex and can be used to elongate the stem regions if desired.

The sequence of the test trigger oligonucleotide is shown in Fig. 20B; this oligonucleotide is termed the alpha primer (SEQ ID NO:25). The alpha primer is complementary to the 3' arm of the T-hairpin as shown in Fig. 20A. When the alpha primer is annealed to the T-hairpin, a cleavage structure is formed that is

recognized by thermostable DNA polymerases. Cleavage of the T-hairpin liberates the 5' single-stranded arm of the T-hairpin, generating the tau primer (SEQ ID NO:26) and a cleaved T-hairpin (Fig. 20B; SEQ ID NO:27). The tau primer is complementary to the 3' arm of the A-hairpin as shown in Fig. 20A. Annealing of the tau primer to the A-hairpin generates another cleavage structure; cleavage of this second cleavage structure liberates the 5' single-stranded arm of the A-hairpin, generating another molecule of the alpha primer which then is annealed to another molecule of the T-hairpin. Thermocycling releases the primers so they can function in additional cleavage reactions. Multiple cycles of annealing and cleavage are carried out. The products of the cleavage reactions are primers and the shortened hairpin structures shown in Fig. 20C. The shortened or cleaved hairpin structures may be resolved from the uncleaved hairpins by electrophoresis on denaturing acrylamide gels.

The annealing and cleavage reactions are carried as follows: In a 50  $\mu$ l reaction volume containing 10 mM Tris-Cl, pH 8.5, 1.0  $\text{MgCl}_2$ , 75 mM KCl, 1 pmole of A-hairpin, 1 pmole T-hairpin, the alpha primer is added at equimolar amount relative to the hairpin structures (1 pmole) or at dilutions ranging from 10- to  $10^6$ -fold and 0.5  $\mu$ l of extract containing a 5' nuclease (prepared as described above) are added. The predicted melting temperature for the alpha or trigger primer is 60°C in the above buffer. Annealing is performed just below this predicted melting temperature at 55°C. Using a Perkin Elmer DNA Thermal Cycler, the reactions are annealed at 55°C for 30 seconds. The temperature is then increased slowly over a five minute period to 72°C to allow for cleavage. After cleavage, the reactions are rapidly brought to 55°C (1°C per second) to allow another cycle of annealing to occur. A range of cycles are performed (20, 40 and 60 cycles) and the reaction products are analyzed at each of these number of cycles. The number of cycles which indicates that the accumulation of cleaved hairpin products has not reached a plateau is then used for subsequent determinations when it is desirable to obtain a quantitative result.

Following the desired number of cycles, the reactions are stopped at 55°C by the addition of 8  $\mu$ l of 95% formamide with 20 mM EDTA and 0.05% marker dyes per 10  $\mu$ l reaction volume. Samples are not heated before loading onto

denaturing polyacrylamide gels (10% polyacrylamide, 19:1 crosslinking, 7 M urea, 89 mM tris-borate, pH 8.3, 2.8 mM EDTA). The samples were not heated to allow for the resolution of single-stranded and re-duplexed uncleaved hairpin molecules.

5 The hairpin molecules may be attached to separate solid support molecules, such as agarose, styrene or magnetic beads, via the 3' end of each hairpin. A spacer molecule may be placed between the 3' end of the hairpin and the bead if so desired. The advantage of attaching the hairpins to a solid support is that this prevents the hybridization of the A- and T-hairpins to one another during the cycles of melting and annealing. The A- and T-hairpins are complementary to one  
10 another (as shown in Fig. 20D) and if allowed to anneal to one another over their entire lengths this would reduce the amount of hairpins available for hybridization to the alpha and tau primers during the detection reaction.

The 5' nucleases of the present invention are used in this assay because they lack significant synthetic activity. The lack of synthetic activity results in the  
15 production of a single cleaved hairpin product (as shown in Fig. 19B, lane 4). Multiple cleavage products may be generated by 1) the presence of interfering synthetic activity (*see* Fig. 19B, lanes 6 and 8) or 2) the presence of primer-independent cleavage in the reaction. The presence of primer-independent cleavage is detected in the trigger/detection assay by the presence of different sized products  
20 at the fork of the cleavage structure. Primer-independent cleavage can be dampened or repressed, when present, by the use of uncleavable nucleotides in the fork region of the hairpin molecule. For example, thiolated nucleotides can be used to replace several nucleotides at the fork region to prevent primer-independent cleavage.

25

## EXAMPLE 5

### Cleavage Of Linear Nucleic Acid Substrates

From the above, it should be clear that native (*i.e.*, "wild type") thermostable DNA polymerases are capable of cleaving hairpin structures in a specific manner and that this discovery can be applied with success to a detection  
30 assay. In this example, the mutant DNAPs of the present invention are tested

against three different cleavage structures shown in Figure 22A. Structure 1 in Figure 22A is simply single stranded 206-mer (the preparation and sequence information for which was discussed above). Structures 2 and 3 are duplexes; structure 2 is the same hairpin structure as shown in Figure 12A (bottom), while structure 3 has the hairpin portion of structure 2 removed.

The cleavage reactions comprised 0.01 pmoles of the resulting substrate DNA, and 1 pmole of pilot oligonucleotide in a total volume of 10  $\mu$ l of 10 mM Tris-Cl, pH 8.3, 100 mM KCl, 1 mM  $MgCl_2$ . Reactions were incubated for 30 minutes at 55°C, and stopped by the addition of 8  $\mu$ l of 95% formamide with 20 mM EDTA and 0.05% marker dyes. Samples were heated to 75°C for 2 minutes immediately before electrophoresis through a 10% polyacrylamide gel (19:1 cross link), with 7M urea, in a buffer of 45 mM Tris-Borate, pH 8.3, 1.4 mM EDTA.


The results were visualized by autoradiography and are shown in Figure 22B with the enzymes indicated as follows: I is native *Taq* DNAP; II is native *Tfl* DNAP; III is Cleavase™ BX shown in Figure 4E; IV is Cleavase™ BB shown in Figure 4F; V is the mutant shown in Figure 5B; and VI is Cleavase™ BN shown in Figure 4G.

Structure 2 was used to "normalize" the comparison. For example, it was found that it took 50 ng of *Taq* DNAP and 300 ng of Cleavase™ BN to give similar amounts of cleavage of Structure 2 in thirty (30) minutes. Under these conditions native *Taq* DNAP is unable to cleave Structure 3 to any significant degree. Native *Tfl* DNAP cleaves Structure 3 in a manner that creates multiple products.

By contrast, all of the mutants tested cleave the linear duplex of Structure 3. This finding indicates that this characteristic of the mutant DNA polymerases is consistent of thermostable polymerases across thermophilic species.

The finding described herein that the mutant DNA polymerases of the present invention are capable of cleaving linear duplex structures allows for application to a more straightforward assay design (Figure 1A). Figure 23 provides a more detailed schematic corresponding to the assay design of Figure 1A.

The two 43-mers depicted in Figure 23 were synthesized by standard methods. Each included a fluorescein on the 5' end for detection purposes and a

biotin on the 3' end to allow attachment to streptavidin coated paramagnetic particles (the biotin-avidin attachment is indicated by "").

Before the trityl groups were removed, the oligos were purified by HPLC to remove truncated by-products of the synthesis reaction. Aliquots of each 43-mer were bound to M-280 Dynabeads (Dynal) at a density of 100 pmoles per mg of beads. Two (2) mgs of beads (200  $\mu$ l) were washed twice in 1X wash/bind buffer (1 M NaCl, 5 mM Tris-Cl, pH 7.5, 0.5 mM EDTA) with 0.1% BSA, 200  $\mu$ l per wash. The beads were magnetically sedimented between washes to allow supernatant removal. After the second wash, the beads were resuspended in 200  $\mu$ l of 2X wash/bind buffer (2 M Na Cl, 10 mM Tris-Cl, pH 7.5 with 1 mM EDTA), and divided into two 100  $\mu$ l aliquots. Each aliquot received 1  $\mu$ l of a 100  $\mu$ M solution of one of the two oligonucleotides. After mixing, the beads were incubated at room temperature for 60 minutes with occasional gentle mixing. The beads were then sedimented and analysis of the supernatants showed only trace amounts of unbound oligonucleotide, indicating successful binding. Each aliquot of beads was washed three times, 100  $\mu$ l per wash, with 1X wash/bind buffer, then twice in a buffer of 10 mM Tris-Cl, pH 8.3 and 75 mM KCl. The beads were resuspended in a final volume of 100  $\mu$ l of the Tris/KCl, for a concentration of 1 pmole of oligo bound to 10  $\mu$ g of beads per  $\mu$ l of suspension. The beads were stored at 4°C between uses.

The types of beads correspond to Figure 1A. That is to say, type 2 beads contain the oligo (SEQ ID NO:33) comprising the complementary sequence (SEQ ID NO:34) for the alpha signal oligo (SEQ ID NO:35) as well as the beta signal oligo (SEQ ID NO:36) which when liberated is a 24-mer. This oligo has no "As" and is "T" rich. Type 3 beads contain the oligo (SEQ ID NO:37) comprising the complementary sequence (SEQ ID NO:38) for the beta signal oligo (SEQ ID NO:39) as well as the alpha signal oligo (SEQ ID NO:35) which when liberated is a 20-mer. This oligo has no "Ts" and is "A" rich.

Cleavage reactions comprised 1  $\mu$ l of the indicated beads, 10 pmoles of unlabelled alpha signal oligo as "pilot" (if indicated) and 500 ng of Cleavase™ BN in 20  $\mu$ l of 75 mM KCl, 10 mM Tris-Cl, pH 8.3, 1.5 mM MgCl<sub>2</sub> and 10  $\mu$ M CTAB. All components except the enzyme were assembled, overlaid with light

mineral oil and warmed to 53°C. The reactions were initiated by the addition of prewarmed enzyme and incubated at that temperature for 30 minutes. Reactions were stopped at temperature by the addition of 16 µl of 95% formamide with 20 mM EDTA and 0.05% each of bromophenol blue and xylene cyanol. This addition  
5 stops the enzyme activity and, upon heating, disrupts the biotin-avidin link, releasing the majority (greater than 95%) of the oligos from the beads. Samples were heated to 75°C for 2 minutes immediately before electrophoresis through a 10% polyacrylamide gel (19:1 cross link), with 7 M urea, in a buffer of 45 mM Tris-Borate, pH 8.3, 1.4 mM EDTA. Results were visualized by contact transfer of  
10 the resolved DNA to positively charged nylon membrane and probing of the blocked membrane with an anti-fluorescein antibody conjugated to alkaline phosphatase. After washing, the signal was developed by incubating the membrane in Western Blue (Promega) which deposits a purple precipitate where the antibody is bound.

15 Figure 24 shows the propagation of cleavage of the linear duplex nucleic acid structures of Figure 23 by the DNAP mutants of the present invention. The two center lanes contain both types of beads. As noted above, the beta signal oligo (SEQ ID NO:36) when liberated is a 24-mer and the alpha signal oligo (SEQ ID NO:35) when liberated is a 20-mer. The formation of the two lower bands  
20 corresponding to the 24-mer and 20-mer is clearly dependent on "pilot".

## EXAMPLE 6

### 5' Exonucleolytic Cleavage ("Nibbling") By Thermostable DNAPs

It has been found that thermostable DNAPs, including those of the present invention, have a true 5' exonuclease capable of nibbling the 5' end of a linear  
25 duplex nucleic acid structures. In this example, the 206 base pair DNA duplex substrate is again employed (see above). In this case, it was produced by the use of one <sup>32</sup>P-labeled primer and one unlabeled primer in a polymerase chain reaction. The cleavage reactions comprised 0.01 pmoles of heat-denatured, end-labeled substrate DNA (with the unlabeled strand also present), 5 pmoles of pilot  
30 oligonucleotide (see pilot oligos in Figure 12A) and 0.5 units of DNAPtaq or 0.5



$\mu$  of Cleavase™ BB in the *E. coli* extract (see above), in a total volume of 10  $\mu$ l of 10 mM Tris•Cl, pH 8.5, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>.

Reactions were initiated at 65°C by the addition of pre-warmed enzyme, then shifted to the final incubation temperature for 30 minutes. The results are shown in Figure 25A. Samples in lanes 1-4 are the results with native *Taq* DNAP, while lanes 5-8 shown the results with Cleavase™ BB. The reactions for lanes 1, 2, 5, and 6 were performed at 65°C and reactions for lanes 3, 4, 7, and 8 were performed at 50°C and all were stopped at temperature by the addition of 8  $\mu$ l of 95% formamide with 20 mM EDTA and 0.05% marker dyes. Samples were heated to 75°C for 2 minutes immediately before electrophoresis through a 10% acrylamide gel (19:1 cross-linked), with 7 M urea, in a buffer of 45 mM Tris•Borate, pH 8.3, 1.4 mM EDTA. The expected product in reactions 1, 2, 5, and 6 is 85 nucleotides long; in reactions 3 and 7, the expected product is 27 nucleotides long. Reactions 4 and 8 were performed without pilot, and should remain at 206 nucleotides. The faint band seen at 24 nucleotides is residual end-labeled primer from the PCR.

The surprising result is that Cleavase™ BB under these conditions causes all of the label to appear in a very small species, suggesting the possibility that the enzyme completely hydrolyzed the substrate. To determine the composition of the fastest-migrating band seen in lanes 5-8 (reactions performed with the deletion mutant), samples of the 206 base pair duplex were treated with either T7 gene 6 exonuclease (USB) or with calf intestine alkaline phosphatase (Promega), according to manufacturers' instructions, to produce either labeled mononucleotide (lane a of Figure 25B) or free <sup>32</sup>P-labeled inorganic phosphate (lane b of Figure 25B), respectively. These products, along with the products seen in lane 7 of panel A were resolved by brief electrophoresis through a 20% acrylamide gel (19:1 cross-link), with 7 M urea, in a buffer of 45 mM Tris•Borate, pH 8.3, 1.4 mM EDTA. Cleavase™ BB is thus capable of converting the substrate to mononucleotides.

**EXAMPLE 7****Nibbling Is Duplex Dependent**

5 The nibbling by Cleavase™ BB is duplex dependent. In this example, internally labeled, single strands of the 206-mer were produced by 15 cycles of primer extension incorporating  $\alpha$ -<sup>32</sup>P labeled dCTP combined with all four unlabeled dNTPs, using an unlabeled 206-bp fragment as a template. Single and double stranded products were resolved by electrophoresis through a non-denaturing 6% polyacrylamide gel (29:1 cross-link) in a buffer of 45 mM Tris•Borate, pH 8.3, 1.4 mM EDTA, visualized by autoradiography, excised from the gel, eluted by  
10 passive diffusion, and concentrated by ethanol precipitation.

The cleavage reactions comprised 0.04 pmoles of substrate DNA, and 2  $\mu$ l of Cleavase™ BB (in an *E. coli* extract as described above) in a total volume of 40  $\mu$ l of 10 mM Tris•Cl, pH 8.5, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>. Reactions were initiated by the addition of pre-warmed enzyme; 10  $\mu$ l aliquots were removed at 5,  
15 10, 20, and 30 minutes, and transferred to prepared tubes containing 8  $\mu$ l of 95% formamide with 30 mM EDTA and 0.05% marker dyes. Samples were heated to 75°C for 2 minutes immediately before electrophoresis through a 10% acrylamide gel (19:1 cross-linked), with 7 M urea, in a buffer of 45 mM Tris•Borate, pH 8.3, 1.4 mM EDTA. Results were visualized by autoradiography as shown in Figure  
20 26. Clearly, the cleavage by Cleavase™ BB depends on a duplex structure; no cleavage of the single strand structure is detected whereas cleavage of the 206-mer duplex is complete.

**EXAMPLE 8****Nibbling Can Be Target Directed**

25 The nibbling activity of the DNAPs of the present invention can be employed with success in a detection assay. One embodiment of such an assay is shown in Figure 27. In this assay, a labelled oligo is employed that is specific for a target sequence. The oligo is in excess of the target so that hybridization is rapid. In this embodiment, the oligo contains two fluorescein labels whose

proximity on the oligo causes their emission to be quenched. When the DNAP is permitted to nibble the oligo the labels separate and are detectable. The shortened duplex is destabilized and disassociates. Importantly, the target is now free to react with an intact labelled oligo. The reaction can continue until the desired level of detection is achieved. An analogous, although different, type of cycling assay has been described employing lambda exonuclease. See C.G. Copley and C. Boot, *BioTechniques* 13:888 (1992).

The success of such an assay depends on specificity. In other words, the oligo must hybridize to the specific target. It is also preferred that the assay be sensitive; the oligo ideally should be able to detect small amounts of target. Figure 28A shows a 5'-end  $^{32}\text{P}$ -labelled primer bound to a plasmid target sequence. In this case, the plasmid was pUC19 (commercially available) which was heat denatured by boiling two (2) minutes and then quick chilling. The primer is a 21-mer (SEQ ID NO:39). The enzyme employed was Cleavase<sup>TM</sup> BX (a dilution equivalent to  $5 \times 10^{-3}$   $\mu\text{l}$  extract) in 100 mM KCl, 10 mM Tris-Cl, pH 8.3, 2 mM  $\text{MnCl}_2$ . The reaction was performed at 55°C for sixteen (16) hours with or without genomic background DNA (from chicken blood). The reaction was stopped by the addition of 8  $\mu\text{l}$  of 95% formamide with 20 mM EDTA and marker dyes.

The products of the reaction were resolved by PAGE (10% polyacrylamide, 19:1 cross link, 1 x TBE) as seen in Figure 28B. Lane "M" contains the labelled 21-mer. Lanes 1-3 contain no specific target, although Lanes 2 and 3 contain 100 ng and 200 ng of genomic DNA, respectively. Lanes 4, 5 and 6 all contain specific target with either 0 ng, 100 ng or 200 ng of genomic DNA, respectively. It is clear that conversion to mononucleotides occurs in Lanes 4, 5 and 6 regardless of the presence or amount of background DNA. Thus, the nibbling can be target directed and specific.

## EXAMPLE 9

### Cleavase Purification

As noted above, expressed thermostable proteins, *i.e.*, the 5' nucleases, were isolated by crude bacterial cell extracts. The precipitated *E. coli* proteins were then, along with other cell debris, removed by centrifugation. In this example, cells expressing the BN clone were cultured and collected (500 grams). For each gram (wet weight) of *E. coli*, 3ml of lysis buffer (50 mM Tris-HCl, pH 8.0, 1 mM EDTA, 100 $\mu$ M NaCl) was added. The cells were lysed with 200 ug/ml lysozyme at room temperature for 20 minutes. Thereafter deoxycholic acid was added to make a 0.2% final concentration and the mixture was incubated 15 minutes at room temperature.

The lysate was sonicated for approximately 6-8 minutes at 0°C. The precipitate was removed by centrifugation (39,000g for 20 minutes). Polyethyleneimine was added (0.5%) to the supernatant and the mixture was incubated on ice for 15 minutes.

The mixture was centrifuged (5,000g for 15 minutes) and the supernatant was retained. This was heated for 30 minutes at 60°C and then centrifuged again (5,000g for 15 minutes) and the supernatant was again retained.

The supernatant was precipitated with 35% ammonium sulfate at 4°C for 15 minutes. The mixture was then centrifuged (5,000g for 15 minutes) and the supernatant was removed. The precipitate was then dissolved in 0.25 M KCl, 20 Tris pH 7.6, 0.2% Tween and 0.1 EDTA) and then dialyzed against Binding Buffer (8X Binding Buffer comprises: 40mM imidazole, 4M NaCl, 160 mM Tris-HCl, pH 7.9).

The solubilized protein is then purified on the Ni<sup>++</sup> column (Novagen). The Binding Buffer is allowed to drain to the top of the column bed and load the column with the prepared extract. A flow rate of about 10 column volumes per hour is optimal for efficient purification. If the flow rate is too fast, more impurities will contaminate the eluted fraction.

The column is washed with 25 ml (10 volumes) of 1X Binding Buffer and then washed with 15 ml (6 volumes) of 1X Wash Buffer (8X Wash Buffer

comprises: 480mM imidazole, 4M NaCl, 160 mM Tris-HCl, pH 7.9). The bound protein was eluted with 15ml (6 volumes) of 1X Elute Buffer (4X Elute Buffer comprises: 4mM imidazole, 2M NaCl, 80 mM Tris-HCl, pH 7.9). Protein is then reprecipitated with 35% Ammonium Sulfate as above. The precipitate was then dissolved and dialyzed against: 20 mM Tris, 100 mM KCl, 1mM EDTA). The solution was brought up to 0.1% each of Tween 20 and NP-40 and stored at 4°C.

From the above, it should be clear that the present invention provides novel cleaving enzymes having heretofore undisclosed nuclease activities. The enzymes can be employed with success in target detection assays of various designs. These assays do not require that the sample DNA be amplified prior to detection and therefore offer an improvement in DNA-based detection technology.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Dahlberg, James E.  
Lyamichev, Victor I.  
Brow, Mary Ann D.
- (ii) TITLE OF INVENTION: 5' Nucleases Derived From Thermostable DNA Polymerase
- (iii) NUMBER OF SEQUENCES: 40
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Haverstock, Medlen & Carroll
  - (B) STREET: 220 Montgomery Street, Suite 2200
  - (C) CITY: San Francisco
  - (D) STATE: California
  - (E) COUNTRY: United States of America
  - (F) ZIP: 94104
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: US
  - (B) FILING DATE: 06-JUN-1994
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 08/073,384
  - (B) FILING DATE: 06-JUN-1993
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 07/986,330
  - (B) FILING DATE: 07-DEC-1992
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Carroll, Peter G.
  - (B) REGISTRATION NUMBER: 32,837
  - (C) REFERENCE/DOCKET NUMBER: FORS-01000
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (415) 705-8410
  - (B) TELEFAX: (415) 397-8338

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 2505 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
 

ATGAGGGGGA TGCTGCCCCT CTTTGAGCCC AAGGGCCGGG TCCTCCTGGT GGACGGCCAC	60
CACCTGGCCT ACCGCACCTT CCACGCCCTG AAGGGCCTCA CCACCAGCCG GGGGGAGCCG	120
GTGCAGGCGG TCTACGGCTT CGCCAAGAGC CTCCTCAAGG CCCTCAAGGA GGACGGGGAC	180



GCGGTGATCG TGGTCTTTGA CGCCAAGGCC CCCTCCTTCC GCCACGAGGC CTACGGGGGG	240
TACAAGGCGG GCCGGGCCCC CACGCCGGAG GACTTTCCCC GGCAACTCGC CCTCATCAAG	300
GAGCTGGTGG ACCTCCTGGG GCTGGCGCGC CTCGAGGTCC CGGGCTACGA GGC GGACGAC	360
GTCTTGCCA GCCTGGCCAA GAAGGCGGAA AAGGAGGGCT ACGAGGTCCG CATCCTCACC	420
GCCGACAAAG ACCTTTACCA GCTCCTTTCC GACCGCATCC ACGTCCTCCA CCCCAGAGGG	480
TACCTCATCA CCCC GGCTG GCTTTGGGAA AAGTACGGCC TGAGGCCCGA CCAGTGGGCC	540
GACTACCGGG CCCTGACCGG GGACGAGTCC GACAACCTTC CCGGGGTCAA GGCATCGGG	600
GAGAAGACGG CGAGGAAGCT TCTGGAGGAG TGGGGGAGCC TGGAAGCCCT CCTCAAGAAC	660
CTGGACCGGC TGAAGCCCGC CATCCGGGAG AAGATCCTGG CCCACATGGA CGATCTGAAG	720
CTCTCCTGGG ACCTGGCCAA GGTGCGCACC GACCTGCCCC TGGAGGTGGA CTTCGCCAAA	780
AGGCGGGAGC CCGACCGGGA GAGGCTTAGG GCCTTTCTGG AGAGGCTTGA GTTTGGCAGC	840
CTCCTCCACG AGTTCGGCCT TCTGGAAAGC CCAAGGCC TGGAGGAGGC CCCCTGGCCC	900
CCGCCGGAAG GGGCCTTCGT GGGCTTTGTG CTTTCCCGCA AGGAGCCCAT GTGGGCCGAT	960
CTTCTGGCCC TGGCCGCCGC CAGGGGGGGC CGGGTCCACC GGGCCCCCGA GCCTTATAAA	1020
GCCCTCAGGG ACCTGAAGGA GGCGCGGGGG CTTCTCGCCA AAGACCTGAG CGTTCTGGCC	1080
CTGAGGGAAG GCCTTGGCCT CCCGCCCGGC GACGACCCA TGCTCCTCGC CTACCTCCTG	1140
GACCCTTCCA ACACCACCCC CGAGGGGGTG GCCCGGCGCT ACGGCGGGGA GTGGACGGAG	1200
GAGGCGGGGG AGCGGGCCGC CCTTTCCGAG AGGCTCTTCG CCAACCTGTG GGGGAGGCTT	1260
GAGGGGGAGG AGAGGCTCCT TTGGCTTTAC CGGGAGGTGG AGAGGCCCTT TTCCGCTGTC	1320
CTGGCCCACA TGGAGGCCAC GGGGGTGCGC CTGGACGTGG CCTATCTCAG GGCCTTGTC	1380
CTGGAGGTGG CCGAGGAGAT CGCCCGCCTC GAGGCCGAGG TCTTCCGCCT GGCCGGCCAC	1440
CCCTTCAACC TCAACTCCCG GGACCAGCTG GAAAGGGTCC TCTTTGACGA GCTAGGGCTT	1500
CCCGCCATCG GCAAGACGGA GAAGACCGGC AAGCGCTCCA CCAGCGCCGC CGTCCTGGAG	1560
GCCCTCCGCG AGGCCACCCC CATCGTGGAG AAGATCCTGC AGTACCGGGA GCTCACCAAG	1620
CTGAAGAGCA CCTACATTGA CCCCTTGCCG GACCTCATCC ACCCCAGGAC GGGCCGCCTC	1680
CACACCCGCT TCAACCAGAC GGCCACGGCC ACGGGCAGGC TAAGTAGCTC CGATCCCAAC	1740
CTCCAGAACA TCCCCGTCCG CACCCCGCTT GGGCAGAGGA TCCGCCGGGC CTTTCATCGCC	1800
GAGGAGGGGT GGCTATTGGT GGCCCTGGAC TATAGCCAGA TAGAGCTCAG GGTGCTGGCC	1860
CACCTCTCCG GCGACGAGAA CCTGATCCGG GTCTTCCAGG AGGGGCGGGA CATCCACACG	1920
GAGACCGCCA GCTGGATGTT CGGCGTCCCC CGGGAGGCCG TGGACCCCTT GATGCGCCGG	1980
GCGGCCAAGA CCATCAACTT CGGGGTCTC TACGGCATGT CGGCCACCG CCTCTCCAG	2040
GAGCTAGCCA TCCTTACGAG GAGGCCAGG CTTTCATTGA GCGCTACTTT CAGAGCTTCC	2100
CCAAGGTGCG GGCCTGGATT GAGAAGACCC TGGAGGAGGG CAGGAGGCGG GGGTACGTGG	2160
AGACCCTCTT CGGCCGCCGC CGCTACGTGC CAGACCTAGA GGCCCGGGTG AAGAGCGTGC	2220

GGGAGGCGGC	CGAGCGCATG	GCCTTCAACA	TGCCCCTCCA	GGGCACCGCC	GCCGACCTCA	2280
TGAAGCTGGC	TATGGTGAAG	CTCTTCCCCA	GGCTGGAGGA	AATGGGGGCC	AGGATGCTCC	2340
TTCAGGTCCA	CGACGAGCTG	GTCCTCGAGG	CCCCAAAAGA	GAGGGCGGAG	GCCGTGGCCC	2400
GGCTGGCCAA	GGAGGTCATG	GAGGGGGTGT	ATCCCCTGGC	CGTGCCCCTG	GAGGTGGAGG	2460
TGGGGATAGG	GGAGGACTGG	CTCTCCGCCA	AGGAGTGATA	CCACC		2505

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2496 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGGCGATGC	TTCCCCTCTT	TGAGCCCAA	GGCCGCGTGC	TCCTGGTGGA	CGGCCACCAC	60
CTGGCCTACC	GCACCTTCTT	TGCCCTCAAG	GGCCTCACCA	CCAGCCGCGG	CGAACCCGTT	120
CAGGCGGTCT	ACGGCTTCGC	CAAAGCCTC	CTCAAGGCC	TGAAGGAGGA	CGGGGACGTG	180
GTGGTGGTGG	TCTTTGACGC	CAAGGCCCC	TCCTTCCGCC	ACGAGGCCTA	CGAGGCCTAC	240
AAGGCGGGCC	GGGCCCCCAC	CCCGGAGGAC	TTTCCCCGGC	AGCTGGCCCT	CATCAAGGAG	300
TTGGTGGACC	TCCTAGGCCT	TGTGCGGCTG	GAGGTTCCCG	GCTTTGAGGC	GGACGACGTG	360
CTGGCCACCC	TGGCCAAGCG	GGCGGAAAAG	GAGGGGTACG	AGGTGCGCAT	CCTCACTGCC	420
GACCGCGACC	TCTACCAGCT	CCTTTCGGAG	CGCATCGCCA	TCCTCCACCC	TGAGGGGTAC	480
CTGATCACCC	CGGCGTGGCT	TTACGAGAAG	TACGGCCTGC	GCCCGGAGCA	GTGGGTGGAC	540
TACCGGGCCC	TGGCGGGGGA	CCCCTCGGAT	AACATCCCCG	GGGTGAAGGG	CATCGGGGAG	600
AAGACCGCCC	AGAGGCTCAT	CCGCGAGTGG	GGGAGCCTGG	AAAACCTCTT	CCAGCACCTG	660
GACCAGGTGA	AGCCCTCCTT	GCGGGAGAAG	CTCCAGGCGG	GCATGGAGGC	CCTGGCCCTT	720
TCCCGGAAGC	TTTCCCAGGT	GCACACTGAC	CTGCCCCTGG	AGGTGGACTT	CGGGAGGCGC	780
CGCACACCCA	ACCTGGAGGG	TCTGCGGGCT	TTTTTGGAGC	GGTTGGAGTT	TGGAAGCCTC	840
CTCCACGAGT	TCGGCCTCCT	GGAGGGGCGG	AAGGCGGCAG	AGGAGGCCCC	CTGGCCCCCT	900
CCGGAAGGGG	CTTTTTTGGG	CTTTTCCTTT	TCCCGTCCCG	AGCCCATGTG	GGCCGAGCTT	960
CTGGCCCTGG	CTGGGGCGTG	GGAGGGGCGC	CTCCATCGGG	CACAAGACCC	CCTTAGGGGC	1020
CTGAGGGACC	TTAAGGGGGT	GCGGGGAATC	CTGGCCAAGG	ACCTGGCGGT	TTTGGCCCTG	1080
CGGGAGGGCC	TGGACCTCTT	CCCAGAGGAC	GACCCCATGC	TCCTGGCCTA	CCTTCTGGAC	1140
CCCTCCAACA	CCACCCCTGA	GGGGGTGGCC	CGGCGTTACG	GGGGGGAGTG	GACGGAGGAT	1200
GCGGGGGAGA	GGGCCCTCCT	GGCCGAGCGC	CTCTTCCAGA	CCCTAAAGGA	GCGCCTTAAG	1260
GGAGAAGAAC	GCCTGCTTTG	GCTTTACGAG	GAGGTGGAGA	AGCCGCTTTC	CCGGGTGTTG	1320
GCCCGGATGG	AGGCCACGGG	GGTCCGGCTG	GACGTGGCCT	ACCTCCAGGC	CCTCTCCCTG	1380

GAGGTGGAGG CGGAGGTGCG CCAGCTGGAG GAGGAGGTCT TCCGCCTGGC CGGCCACCCC	1440
TTCAACCTCA ACTCCCGCGA CCAGCTGGAG CGGGTGCTCT TTGACGAGCT GGGCCTGCCT	1500
GCCATCGGCA AGACGGAGAA GACGGGGAAA CGCTCCACCA GCGCTGCCGT GCTGGAGGCC	1560
CTGCGAGAGG CCCACCCCAT CGTGGACCGC ATCCTGCAGT ACCGGGAGCT CACCAAGCTC	1620
AAGAACACCT ACATAGACCC CCTGCCCCGCC CTGGTCCACC CCAAGACCGG CCGGCTCCAC	1680
ACCCGCTTCA ACCAGACGGC CACCGCCACG GGCAGGCTTT CCAGCTCCGA CCCCAACCTG	1740
CAGAACATCC CCGTGCGCAC CCCTCTGGGC CAGCGCATCC GCCGAGCCTT CGTGGCCGAG	1800
GAGGGCTGGG TGCTGGTGGT CTTGGACTAC AGCCAGATTG AGCTTCGGGT CCTGGCCCAC	1860
CTCTCCGGGG ACGAGAACCT GATCCGGGTC TTTCAGGAGG GGAGGGACAT CCACACCCAG	1920
ACCGCCAGCT GGATGTTCGG CGTTTCCCCC GAAGGGGTAG ACCCTCTGAT GCGCCGGGCG	1980
GCCAAGACCA TCAACTTCGG GGTGCTCTAC GGCATGTCCG CCCACCGCCT CTCCGGGGAG	2040
CTTTCCATCC CCTACGAGGA GCGGGTGGCC TTCATTGAGC GCTACTTCCA GAGCTACCCC	2100
AAGGTGCGGG CCTGGATTGA GGGGACCCTC GAGGAGGGCC GCCGGCGGGG GTATGTGGAG	2160
ACCCTCTTCG GCCGCCGGCG CTATGTGCCC GACCTCAACG CCCGGGTGAA GAGCGTGCGC	2220
GAGGCGGCGG AGCGCATGGC CTTCAACATG CCGGTCCAGG GCACCGCCGC CGACCTCATG	2280
AAGCTGGCCA TGGTGCGGCT TTTCCCCCGG CTTCAGGAAC TGGGGGCGAG GATGCTTTTG	2340
CAGGTGCACG ACGAGCTGGT CCTCGAGGCC CCCAAGGACC GGGCGGAGAG GGTAGCCGCT	2400
TTGGCCAAGG AGGTCATGGA GGGGGTCTGG CCCCTGCAGG TGCCCCTGGA GGTGGAGGTG	2460
GGCCTGGGGG AGGACTGGCT CTCCGCCAAG GAGTAG	2496

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2504 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATGGAGGCGA TGCTTCCGCT CTTTGAACCC AAAGGCCGGG TCCTCCTGGT GGACGGCCAC	60
CACCTGGCCT ACCGCACCTT CTTGCCCCTG AAGGGCCTCA CCACGAGCCG GGGCGAACCG	120
GTGCAGGCGG TCTACGGCTT CGCCAAGAGC CTCCTCAAGG CCCTGAAGGA GGACGGGTAC	180
AAGGCCGTCT TCGTGGTCTT TGACGCCAAG GCCCCCTCCT TCCGCCACGA GGCCTACGAG	240
GCCTACAAGG CGGGGAGGGC CCCGACCCCC GAGGACTTCC CCCGGCAGCT CGCCCTCATC	300
AAGGAGCTGG TGGACCTCCT GGGGTTTACC CGCCTCGAGG TCCCCGGCTA CGAGGCGGAC	360
GACGTTCTCG CCACCCTGGC CAAGAAGGCG GAAAAGGAGG GGTACGAGGT GCGCATCCTC	420
ACCGCCGACC GCGACCTCTA CCAACTCGTC TCCGACCGCG TCGCCGTCCT CCACCCCGAG	480
GGCCACCTCA TCACCCCGGA GTGGCTTTGG GAGAAGTACG GCCTCAGGCC GGAGCAGTGG	540

GTGGACTTCC	GCGCCCTCGT	GGGGGACCCC	TCCGACAACC	TCCCCGGGGT	CAAGGGCATC	600
GGGGAGAAGA	CCGCCCTCAA	GCTCCTCAAG	GAGTGGGGAA	GCCTGGAAAA	CCTCCTCAAG	660
AACCTGGACC	GGGTAAAGCC	AGAAAACGTC	CGGGAGAAGA	TCAAGGCCCA	CCTGGAAGAC	720
CTCAGGCTCT	CCTTGGAGCT	CTCCCGGGTG	CGCACCGACC	TCCCCCTGGA	GGTGGACCTC	780
GCCCAGGGGC	GGGAGCCCGA	CCGGGAGGGG	CTTAGGGCCT	TCCTGGAGAG	GCTGGAGTTC	840
GGCAGCCTCC	TCCACGAGTT	CGGCCTCCTG	GAGGCCCCCG	CCCCCCTGGA	GGAGGCCCCC	900
TGGCCCCCGC	CGGAAGGGGC	CTTCGTGGGC	TTCGTCTCT	CCCGCCCCGA	GCCCATGTGG	960
GCGGAGCTTA	AAGCCCTGGC	CGCCTGCAGG	GACGGCCGGG	TGCACCGGGC	AGCAGACCCC	1020
TTGGCGGGGC	TAAAGGACCT	CAAGGAGGTC	CGGGGCCTCC	TCGCCAAGGA	CCTCGCCGTC	1080
TTGGCCTCGA	GGGAGGGGCT	AGACCTCGTG	CCCGGGGACG	ACCCCATGCT	CCTCGCCTAC	1140
CTCCTGGACC	CCTCCAACAC	CACCCCCGAG	GGGGTGGCGC	GGCGCTACGG	GGGGGAGTGG	1200
ACGGAGGACG	CCGCCCACCG	GGCCCTCCTC	TCGGAGAGGC	TCCATCGGAA	CCTCCTTAAG	1260
CGCCTCGAGG	GGGAGGAGAA	GCTCCTTTGG	CTCTACCACG	AGGTGGAAAA	GCCCCTCTCC	1320
CGGGTCCTGG	CCCACATGGA	GGCCACCGGG	GTACGGCTGG	ACGTGGCCTA	CCTTCAGGCC	1380
CTTTCCCTGG	AGCTTGCGGA	GGAGATCCGC	CGCCTCGAGG	AGGAGGTCTT	CCGCTTGGCG	1440
GGCCACCCCT	TCAACCTCAA	CTCCCGGGAC	CAGCTGGAAA	GGGTGCTCTT	TGACGAGCTT	1500
AGGCTTCCCG	CCTTGGGGAA	GACGCAAAAG	ACAGGCAAGC	GCTCCACCAG	CGCCGCGGTG	1560
CTGGAGGCCC	TACGGGAGGC	CCACCCCATC	GTGGAGAAGA	TCCTCCAGCA	CCGGGAGCTC	1620
ACCAAGCTCA	AGAACACCTA	CGTGGACCCC	CTCCCAAGCC	TCGTCCACCC	GAGGACGGGC	1680
CGCCTCCACA	CCCGCTTCAA	CCAGACGGCC	ACGGCCACGG	GGAGGCTTAG	TAGCTCCGAC	1740
CCCAACCTGC	AGAACATCCC	CGTCCGCACC	CCCTTGGGCC	AGAGGATCCG	CCGGGCCTTC	1800
GTGGCCGAGG	CGGGTTGGGC	GTTGGTGGCC	CTGGACTATA	GCCAGATAGA	GCTCCGCGTC	1860
CTCGCCCACC	TCTCCGGGGA	CGAAAACCTG	ATCAGGGTCT	TCCAGGAGGG	GAAGGACATC	1920
CACACCCAGA	CCGCAAGCTG	GATGTTTCGGC	GTCCCCCGGG	AGGCCGTGGA	CCCCCTGATG	1980
CGCCGGGCGG	CCAAGACGGT	GAACCTTCGGC	GTCCTCTACG	GCATGTCGCGC	CCATAGGCTC	2040
TCCCAGGAGC	TTGCCATCCC	CTACGAGGAG	GCGGTGGCCT	TTATAGAGGC	TACTTCCAAA	2100
GCTTCCCCAA	GGTGCGGGCC	TGGATAGAAA	AGACCCTGGA	GGAGGGGAGG	AAGCGGGGCT	2160
ACGTGGAAAC	CCTCTTCGGA	AGAAGGCGCT	ACGTGCCCGA	CCTCAACGCC	CGGGTGAAGA	2220
GCGTCAGGGA	GGCCGCGGAG	CGCATGGCCT	TCAACATGCC	CGTCCAGGGC	ACCGCCGCCG	2280
ACCTCATGAA	GCTCGCCATG	GTGAAGCTCT	TCCCCCGCCT	CCGGGAGATG	GGGGCCCCGA	2340
TGCTCCTCCA	GGTCCACGAC	GAGCTCCTCC	TGGAGGCCCC	CCAAGCGCGG	GCCGAGGAGG	2400
TGGCGGCTTT	GGCCAAGGAG	GCCATGGAGA	AGGCCTATCC	CCTCGCCGTG	CCCCTGGAGG	2460
TGGAGGTGGG	GATGGGGGAG	GACTGGCTTT	CCGCCAAGGG	TTAG		2504

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 832 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Arg Gly Met Leu Pro Leu Phe Glu Pro Lys Gly Arg Val Leu Leu
 1           5           10           15
Val Asp Gly His His Leu Ala Tyr Arg Thr Phe His Ala Leu Lys Gly
          20           25           30
Leu Thr Thr Ser Arg Gly Glu Pro Val Gln Ala Val Tyr Gly Phe Ala
          35           40           45
Lys Ser Leu Leu Lys Ala Leu Lys Glu Asp Gly Asp Ala Val Ile Val
          50           55           60
Val Phe Asp Ala Lys Ala Pro Ser Phe Arg His Glu Ala Tyr Gly Gly
          65           70           75           80
Tyr Lys Ala Gly Arg Ala Pro Thr Pro Glu Asp Phe Pro Arg Gln Leu
          85           90           95
Ala Leu Ile Lys Glu Leu Val Asp Leu Leu Gly Leu Ala Arg Leu Glu
          100          105          110
Val Pro Gly Tyr Glu Ala Asp Asp Val Leu Ala Ser Leu Ala Lys Lys
          115          120          125
Ala Glu Lys Glu Gly Tyr Glu Val Arg Ile Leu Thr Ala Asp Lys Asp
          130          135          140
Leu Tyr Gln Leu Leu Ser Asp Arg Ile His Val Leu His Pro Glu Gly
          145          150          155          160
Tyr Leu Ile Thr Pro Ala Trp Leu Trp Glu Lys Tyr Gly Leu Arg Pro
          165          170          175
Asp Gln Trp Ala Asp Tyr Arg Ala Leu Thr Gly Asp Glu Ser Asp Asn
          180          185          190
Leu Pro Gly Val Lys Gly Ile Gly Glu Lys Thr Ala Arg Lys Leu Leu
          195          200          205
Glu Glu Trp Gly Ser Leu Glu Ala Leu Leu Lys Asn Leu Asp Arg Leu
          210          215          220
Lys Pro Ala Ile Arg Glu Lys Ile Leu Ala His Met Asp Asp Leu Lys
          225          230          235          240
Leu Ser Trp Asp Leu Ala Lys Val Arg Thr Asp Leu Pro Leu Glu Val
          245          250          255
Asp Phe Ala Lys Arg Arg Glu Pro Asp Arg Glu Arg Leu Arg Ala Phe
          260          265          270
Leu Glu Arg Leu Glu Phe Gly Ser Leu Leu His Glu Phe Gly Leu Leu
          275          280          285

```

Glu	Ser	Pro	Lys	Ala	Leu	Glu	Glu	Ala	Pro	Trp	Pro	Pro	Pro	Glu	Gly			
	290					295					300							
Ala	Phe	Val	Gly	Phe	Val	Leu	Ser	Arg	Lys	Glu	Pro	Met	Trp	Ala	Asp			
305					310					315					320			
Leu	Leu	Ala	Leu	Ala	Ala	Ala	Arg	Gly	Gly	Arg	Val	His	Arg	Ala	Pro			
				325					330					335				
Glu	Pro	Tyr	Lys	Ala	Leu	Arg	Asp	Leu	Lys	Glu	Ala	Arg	Gly	Leu	Leu			
			340					345					350					
Ala	Lys	Asp	Leu	Ser	Val	Leu	Ala	Leu	Arg	Glu	Gly	Leu	Gly	Leu	Pro			
		355					360					365						
Pro	Gly	Asp	Asp	Pro	Met	Leu	Leu	Ala	Tyr	Leu	Leu	Asp	Pro	Ser	Asn			
	370					375					380							
Thr	Thr	Pro	Glu	Gly	Val	Ala	Arg	Arg	Tyr	Gly	Gly	Glu	Trp	Thr	Glu			
385					390					395					400			
Glu	Ala	Gly	Glu	Arg	Ala	Ala	Leu	Ser	Glu	Arg	Leu	Phe	Ala	Asn	Leu			
				405					410					415				
Trp	Gly	Arg	Leu	Glu	Gly	Glu	Glu	Arg	Leu	Leu	Trp	Leu	Tyr	Arg	Glu			
			420					425					430					
Val	Glu	Arg	Pro	Leu	Ser	Ala	Val	Leu	Ala	His	Met	Glu	Ala	Thr	Gly			
		435					440					445						
Val	Arg	Leu	Asp	Val	Ala	Tyr	Leu	Arg	Ala	Leu	Ser	Leu	Glu	Val	Ala			
	450					455					460							
Glu	Glu	Ile	Ala	Arg	Leu	Glu	Ala	Glu	Val	Phe	Arg	Leu	Ala	Gly	His			
465					470					475					480			
Pro	Phe	Asn	Leu	Asn	Ser	Arg	Asp	Gln	Leu	Glu	Arg	Val	Leu	Phe	Asp			
				485					490					495				
Glu	Leu	Gly	Leu	Pro	Ala	Ile	Gly	Lys	Thr	Glu	Lys	Thr	Gly	Lys	Arg			
			500					505					510					
Ser	Thr	Ser	Ala	Ala	Val	Leu	Glu	Ala	Leu	Arg	Glu	Ala	His	Pro	Ile			
		515					520					525						
Val	Glu	Lys	Ile	Leu	Gln	Tyr	Arg	Glu	Leu	Thr	Lys	Leu	Lys	Ser	Thr			
	530					535					540							
Tyr	Ile	Asp	Pro	Leu	Pro	Asp	Leu	Ile	His	Pro	Arg	Thr	Gly	Arg	Leu			
545					550					555					560			
His	Thr	Arg	Phe	Asn	Gln	Thr	Ala	Thr	Ala	Thr	Gly	Arg	Leu	Ser	Ser			
				565					570					575				
Ser	Asp	Pro	Asn	Leu	Gln	Asn	Ile	Pro	Val	Arg	Thr	Pro	Leu	Gly	Gln			
			580					585					590					
Arg	Ile	Arg	Arg	Ala	Phe	Ile	Ala	Glu	Glu	Gly	Trp	Leu	Leu	Val	Ala			
		595					600					605						
Leu	Asp	Tyr	Ser	Gln	Ile	Glu	Leu	Arg	Val	Leu	Ala	His	Leu	Ser	Gly			
	610					615					620							
Asp	Glu	Asn	Leu	Ile	Arg	Val	Phe	Gln	Glu	Gly	Arg	Asp	Ile	His	Thr			
625					630					635					640			



(2) INFORMATION FOR SEQ ID NO:5:

(A) LENGTH: 831 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

-83-

Leu Ile Lys Glu Leu Val Asp Leu Leu Gly Leu Val Arg Leu Glu Val  
 100 105 110  
 Pro Gly Phe Glu Ala Asp Asp Val Leu Ala Thr Leu Ala Lys Arg Ala  
 115 120 125  
 Glu Lys Glu Gly Tyr Glu Val Arg Ile Leu Thr Ala Asp Arg Asp Leu  
 130 135 140  
 Tyr Gln Leu Leu Ser Glu Arg Ile Ala Ile Leu His Pro Glu Gly Tyr  
 145 150 155 160  
 Leu Ile Thr Pro Ala Trp Leu Tyr Glu Lys Tyr Gly Leu Arg Pro Glu  
 165 170 175  
 Gln Trp Val Asp Tyr Arg Ala Leu Ala Gly Asp Pro Ser Asp Asn Ile  
 180 185 190  
 Pro Gly Val Lys Gly Ile Gly Glu Lys Thr Ala Gln Arg Leu Ile Arg  
 195 200 205  
 Glu Trp Gly Ser Leu Glu Asn Leu Phe Gln His Leu Asp Gln Val Lys  
 210 215 220  
 Pro Ser Leu Arg Glu Lys Leu Gln Ala Gly Met Glu Ala Leu Ala Leu  
 225 230 235 240  
 Ser Arg Lys Leu Ser Gln Val His Thr Asp Leu Pro Leu Glu Val Asp  
 245 250 255  
 Phe Gly Arg Arg Arg Thr Pro Asn Leu Glu Gly Leu Arg Ala Phe Leu  
 260 265 270  
 Glu Arg Leu Glu Phe Gly Ser Leu Leu His Glu Phe Gly Leu Leu Glu  
 275 280 285  
 Gly Pro Lys Ala Ala Glu Glu Ala Pro Trp Pro Pro Pro Glu Gly Ala  
 290 295 300  
 Phe Leu Gly Phe Ser Phe Ser Arg Pro Glu Pro Met Trp Ala Glu Leu  
 305 310 315 320  
 Leu Ala Leu Ala Gly Ala Trp Glu Gly Arg Leu His Arg Ala Gln Asp  
 325 330 335  
 Pro Leu Arg Gly Leu Arg Asp Leu Lys Gly Val Arg Gly Ile Leu Ala  
 340 345 350  
 Lys Asp Leu Ala Val Leu Ala Leu Arg Glu Gly Leu Asp Leu Phe Pro  
 355 360 365  
 Glu Asp Asp Pro Met Leu Leu Ala Tyr Leu Leu Asp Pro Ser Asn Thr  
 370 375 380  
 Thr Pro Glu Gly Val Ala Arg Arg Tyr Gly Gly Glu Trp Thr Glu Asp  
 385 390 395 400  
 Ala Gly Glu Arg Ala Leu Leu Ala Glu Arg Leu Phe Gln Thr Leu Lys  
 405 410 415  
 Glu Arg Leu Lys Gly Glu Glu Arg Leu Leu Trp Leu Tyr Glu Glu Val  
 420 425 430  
 Glu Lys Pro Leu Ser Arg Val Leu Ala Arg Met Glu Ala Thr Gly Val  
 435 440 445

Arg	Leu	Asp	Val	Ala	Tyr	Leu	Gln	Ala	Leu	Ser	Leu	Glu	Val	Glu	Ala
450						455					460				
Glu	Val	Arg	Gln	Leu	Glu	Glu	Glu	Val	Phe	Arg	Leu	Ala	Gly	His	Pro
465					470					475					480
Phe	Asn	Leu	Asn	Ser	Arg	Asp	Gln	Leu	Glu	Arg	Val	Leu	Phe	Asp	Glu
				485					490					495	
Leu	Gly	Leu	Pro	Ala	Ile	Gly	Lys	Thr	Glu	Lys	Thr	Gly	Lys	Arg	Ser
			500					505					510		
Thr	Ser	Ala	Ala	Val	Leu	Glu	Ala	Leu	Arg	Glu	Ala	His	Pro	Ile	Val
		515					520					525			
Asp	Arg	Ile	Leu	Gln	Tyr	Arg	Glu	Leu	Thr	Lys	Leu	Lys	Asn	Thr	Tyr
	530					535					540				
Ile	Asp	Pro	Leu	Pro	Ala	Leu	Val	His	Pro	Lys	Thr	Gly	Arg	Leu	His
545					550					555					560
Thr	Arg	Phe	Asn	Gln	Thr	Ala	Thr	Ala	Thr	Gly	Arg	Leu	Ser	Ser	Ser
				565					570					575	
Asp	Pro	Asn	Leu	Gln	Asn	Ile	Pro	Val	Arg	Thr	Pro	Leu	Gly	Gln	Arg
			580					585					590		
Ile	Arg	Arg	Ala	Phe	Val	Ala	Glu	Glu	Gly	Trp	Val	Leu	Val	Val	Leu
		595					600					605			
Asp	Tyr	Ser	Gln	Ile	Glu	Leu	Arg	Val	Leu	Ala	His	Leu	Ser	Gly	Asp
	610					615					620				
Glu	Asn	Leu	Ile	Arg	Val	Phe	Gln	Glu	Gly	Arg	Asp	Ile	His	Thr	Gln
625					630					635					640
Thr	Ala	Ser	Trp	Met	Phe	Gly	Val	Ser	Pro	Glu	Gly	Val	Asp	Pro	Leu
				645					650					655	
Met	Arg	Arg	Ala	Ala	Lys	Thr	Ile	Asn	Phe	Gly	Val	Leu	Tyr	Gly	Met
			660					665					670		
Ser	Ala	His	Arg	Leu	Ser	Gly	Glu	Leu	Ser	Ile	Pro	Tyr	Glu	Glu	Ala
		675					680					685			
Val	Ala	Phe	Ile	Glu	Arg	Tyr	Phe	Gln	Ser	Tyr	Pro	Lys	Val	Arg	Ala
	690					695					700				
Trp	Ile	Glu	Gly	Thr	Leu	Glu	Glu	Gly	Arg	Arg	Arg	Gly	Tyr	Val	Glu
705					710					715					720
Thr	Leu	Phe	Gly	Arg	Arg	Arg	Tyr	Val	Pro	Asp	Leu	Asn	Ala	Arg	Val
				725					730					735	
Lys	Ser	Val	Arg	Glu	Ala	Ala	Glu	Arg	Met	Ala	Phe	Asn	Met	Pro	Val
			740					745					750		
Gln	Gly	Thr	Ala	Ala	Asp	Leu	Met	Lys	Leu	Ala	Met	Val	Arg	Leu	Phe
		755					760					765			
Pro	Arg	Leu	Gln	Glu	Leu	Gly	Ala	Arg	Met	Leu	Leu	Gln	Val	His	Asp
	770					775					780				
Glu	Leu	Val	Leu	Glu	Ala	Pro	Lys	Asp	Arg	Ala	Glu	Arg	Val	Ala	Ala
785					790					795					800

(2) INFORMATION FOR SEQ ID NO:6:

(A) LENGTH: 834 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

-86-

Glu Val Asp Leu Ala Gln Gly Arg Glu Pro Asp Arg Glu Gly Leu Arg  
 260 265 270  
 Ala Phe Leu Glu Arg Leu Glu Phe Gly Ser Leu Leu His Glu Phe Gly  
 275 280 285  
 Leu Leu Glu Ala Pro Ala Pro Leu Glu Glu Ala Pro Trp Pro Pro Pro  
 290 295 300  
 Glu Gly Ala Phe Val Gly Phe Val Leu Ser Arg Pro Glu Pro Met Trp  
 305 310 315 320  
 Ala Glu Leu Lys Ala Leu Ala Ala Cys Arg Asp Gly Arg Val His Arg  
 325 330 335  
 Ala Ala Asp Pro Leu Ala Gly Leu Lys Asp Leu Lys Glu Val Arg Gly  
 340 345 350  
 Leu Leu Ala Lys Asp Leu Ala Val Leu Ala Ser Arg Glu Gly Leu Asp  
 355 360 365  
 Leu Val Pro Gly Asp Asp Pro Met Leu Leu Ala Tyr Leu Leu Asp Pro  
 370 375 380  
 Ser Asn Thr Thr Pro Glu Gly Val Ala Arg Arg Tyr Gly Gly Glu Trp  
 385 390 395 400  
 Thr Glu Asp Ala Ala His Arg Ala Leu Leu Ser Glu Arg Leu His Arg  
 405 410 415  
 Asn Leu Leu Lys Arg Leu Glu Gly Glu Glu Lys Leu Leu Trp Leu Tyr  
 420 425 430  
 His Glu Val Glu Lys Pro Leu Ser Arg Val Leu Ala His Met Glu Ala  
 435 440 445  
 Thr Gly Val Arg Leu Asp Val Ala Tyr Leu Gln Ala Leu Ser Leu Glu  
 450 455 460  
 Leu Ala Glu Glu Ile Arg Arg Leu Glu Glu Glu Val Phe Arg Leu Ala  
 465 470 475 480  
 Gly His Pro Phe Asn Leu Asn Ser Arg Asp Gln Leu Glu Arg Val Leu  
 485 490 495  
 Phe Asp Glu Leu Arg Leu Pro Ala Leu Gly Lys Thr Gln Lys Thr Gly  
 500 505 510  
 Lys Arg Ser Thr Ser Ala Ala Val Leu Glu Ala Leu Arg Glu Ala His  
 515 520 525  
 Pro Ile Val Glu Lys Ile Leu Gln His Arg Glu Leu Thr Lys Leu Lys  
 530 535 540  
 Asn Thr Tyr Val Asp Pro Leu Pro Ser Leu Val His Pro Arg Thr Gly  
 545 550 555 560  
 Arg Leu His Thr Arg Phe Asn Gln Thr Ala Thr Ala Thr Gly Arg Leu  
 565 570 575  
 Ser Ser Ser Asp Pro Asn Leu Gln Asn Ile Pro Val Arg Thr Pro Leu  
 580 585 590  
 Gly Gln Arg Ile Arg Arg Ala Phe Val Ala Glu Ala Gly Trp Ala Leu  
 595 600 605

Val Ala Leu Asp Tyr Ser Gln Ile Glu Leu Arg Val Leu Ala His Leu  
 610 615 620  
 Ser Gly Asp Glu Asn Leu Ile Arg Val Phe Gln Glu Gly Lys Asp Ile  
 625 630 635 640  
 His Thr Gln Thr Ala Ser Trp Met Phe Gly Val Pro Pro Glu Ala Val  
 645 650 655  
 Asp Pro Leu Met Arg Arg Ala Ala Lys Thr Val Asn Phe Gly Val Leu  
 660 665 670  
 Tyr Gly Met Ser Ala His Arg Leu Ser Gln Glu Leu Ala Ile Pro Tyr  
 675 680 685  
 Glu Glu Ala Val Ala Phe Ile Glu Arg Tyr Phe Gln Ser Phe Pro Lys  
 690 695 700  
 Val Arg Ala Trp Ile Glu Lys Thr Leu Glu Glu Gly Arg Lys Arg Gly  
 705 710 715 720  
 Tyr Val Glu Thr Leu Phe Gly Arg Arg Arg Tyr Val Pro Asp Leu Asn  
 725 730 735  
 Ala Arg Val Lys Ser Val Arg Glu Ala Ala Glu Arg Met Ala Phe Asn  
 740 745 750  
 Met Pro Val Gln Gly Thr Ala Ala Asp Leu Met Lys Leu Ala Met Val  
 755 760 765  
 Lys Leu Phe Pro Arg Leu Arg Glu Met Gly Ala Arg Met Leu Leu Gln  
 770 775 780  
 Val His Asp Glu Leu Leu Leu Glu Ala Pro Gln Ala Arg Ala Glu Glu  
 785 790 795 800  
 Val Ala Ala Leu Ala Lys Glu Ala Met Glu Lys Ala Tyr Pro Leu Ala  
 805 810 815  
 Val Pro Leu Glu Val Glu Val Gly Met Gly Glu Asp Trp Leu Ser Ala  
 820 825 830  
 Lys Gly

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2502 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATGNNGGCGA TGCTTCCCCT CTTTGAGCCC AAAGGCCGGG TCCTCCTGGT GGACGGCCAC	60
CACCTGGCCT ACCGCACCTT CTTGCGCCTG AAGGGCCTCA CCACCAGCCG GGGCGAACCG	120
GTGCAGGCGG TCTACGGCTT CGCCAAGAGC CTCCTCAAGG CCCTGAAGGA GGACGGGGAC	180
NNGGCGGTGN TCGTGGTCTT TGACGCCAAG GCCCCCTCCT TCCGCCACGA GGCCTACGAG	240
GCCTACAAGG CGGGCCGGGC CCCACCCCG GAGGACTTTC CCCGGCAGCT CGCCCTCATC	300



AAGGAGCTGG TGGACCTCCT GGGGCTTGCG CGCCTCGAGG TCCCCGGCTA CGAGGCGGAC	360
GACGTNCTGG CCACCCTGGC CAAGAAGGCG GAAAAGGAGG GGTACGAGGT GCGCATCCTC	420
ACCGCCGACC GCGACCTCTA CCAGCTCCTT TCCGACCGCA TCGCCGTCCT CCACCCCGAG	480
GGGTACCTCA TCACCCCGGC GTGGCTTTGG GAGAAGTACG GCCTGAGGCC GGAGCAGTGG	540
GTGGACTACC GGGCCCTGGC GGGGGACCCC TCCGACAACC TCCCCGGGGT CAAGGGCATC	600
GGGGAGAAGA CCGCCCNAA GCTCCTCNAG GAGTGGGGGA GCCTGGAAAA CCTCCTCAAG	660
AACCTGGACC GGGTGAAGCC CGCCNTCCGG GAGAAGATCC AGGCCACAT GGANGACCTG	720
ANGCTCTCCT GGGAGCTNTC CCAGGTGCGC ACCGACCTGC CCCTGGAGGT GGACTTCGCC	780
AAGNGGCGGG AGCCCGACCG GGAGGGGCTT AGGGCCTTTC TGGAGAGGCT GGAGTTTGGC	840
AGCCTCCTCC ACGAGTTCGG CCTCCTGGAG GGCCCCAAGG CCCTGGAGGA GGCCCCCTGG	900
CCCCCGCCGG AAGGGGCCTT CGTGGGCTTT GTCCTTTCCC GCCCCGAGCC CATGTGGGCC	960
GAGCTTCTGG CCCTGGCCGC CGCCAGGGAG GGCCGGGTCC ACCGGGCACC AGACCCCTTT	1020
ANGGGCCTNA GGGACCTNAA GGAGGTGCGG GGNCTCCTCG CCAAGGACCT GGCCGTTTTG	1080
GCCCTGAGGG AGGGCCTNGA CCTCNTGCCC GGGGACGACC CCATGCTCCT CGCCTACCTC	1140
CTGGACCCCT CCAACACCAC CCCCAGAGGG GTGGCCCGGC GCTACGGGGG GGAGTGGACG	1200
GAGGANGCGG GGGAGCGGGC CCTCCTNTCC GAGAGGCTCT TCCNGAACCT NNNGCAGCGC	1260
CTTGAGGGGG AGGAGAGGCT CCTTTGGCTT TACCAGGAGG TGGAGAAGCC CCTTTCCCGG	1320
GTCCTGGCCC ACATGGAGGC CACGGGGGTN CGGCTGGACG TGGCCTACCT CCAGGCCCTN	1380
TCCCTGGAGG TGGCGGAGGA GATCCGCCGC CTCGAGGAGG AGGTCTTCCG CCTGGCCGGC	1440
CACCCCTTCA ACCTCAACTC CCGGGACCAG CTGGAAAGGG TGCTCTTTGA CGAGCTNGGG	1500
CTTCCCGCCA TCGGCAAGAC GGAGAAGACN GGCAAGCGCT CCACCAGCGC CGCCGTGCTG	1560
GAGGCCCTNC GNGAGGCCCA CCCCATCGTG GAGAAGATCC TGCAGTACCG GGAGCTCACC	1620
AAGCTCAAGA ACACCTACAT NGACCCCTG CCNGNCCTCG TCCACCCAG GACGGGCCGC	1680
CTCCACACCC GCTTCAACCA GACGGCCACG GCCACGGGCA GGCTTAGTAG CTCCGACCCC	1740
AACCTGCAGA ACATCCCCGT CCGCACCCCN CTGGGCCAGA GGATCCGCCG GGCCTTCGTG	1800
GCCGAGGAGG GNTGGGTGTT GGTGGCCCTG GACTATAGCC AGATAGAGCT CCGGGTCCTG	1860
GCCCACCTCT CCGGGGACGA GAACCTGATC CGGGTCTTCC AGGAGGGGAG GGACATCCAC	1920
ACCCAGACCG CCAGCTGGAT GTTCGGCGTC CCCCCGAGG CCGTGGACCC CCTGATGCGC	1980
CGGGCGGCCA AGACCATCAA CTTCGGGGTC CTCTACGGCA TGTCCGCCCA CCGCCTCTCC	2040
CAGGAGCTTG CCATCCCCTA CGAGGAGGCG GTGGCCTTCA TTGAGCGCTA CTTCCAGAGC	2100
TTCCCCAAGG TCGGGGCTG GATTGAGAAG ACCCTGGAGG AGGGCAGGAG GCGGGGGTAC	2160
GTGGAGACCC TCTTCGGCCG CCGGCGCTAC GTGCCCGACC TCAACGCCCG GGTGAAGAGC	2220
GTGCGGGAGG CGGCGGAGCG CATGGCCTTC AACATGCCCG TCCAGGGCAC CGCCGCCGAC	2280
CTCATGAAGC TGGCCATGGT GAAGCTCTTC CCCCCGCTNC AGGAAATGGG GGCCAGGATG	2340

CTCCTNCAGG TCCACGACGA GCTGGTCCTC GAGGCCCCCA AAGAGCGGGC GGAGGNGGTG 2400  
 GCCGCTTTGG CCAAGGAGGT CATGGAGGGG GTCTATCCCC TGGCCGTGCC CCTGGAGGTG 2460  
 GAGGTGGGGA TGGGGGAGGA CTGGCTCTCC GCCAAGGAGT AG 2502

## (2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 833 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met	Xaa	Ala	Met	Leu	Pro	Leu	Phe	Glu	Pro	Lys	Gly	Arg	Val	Leu	Leu	1	5	10	15
Val	Asp	Gly	His	His	Leu	Ala	Tyr	Arg	Thr	Phe	Phe	Ala	Leu	Lys	Gly	20	25	30	
Leu	Thr	Thr	Ser	Arg	Gly	Glu	Pro	Val	Gln	Ala	Val	Tyr	Gly	Phe	Ala	35	40	45	
Lys	Ser	Leu	Leu	Lys	Ala	Leu	Lys	Glu	Asp	Gly	Asp	Ala	Val	Xaa	Val	50	55	60	
Val	Phe	Asp	Ala	Lys	Ala	Pro	Ser	Phe	Arg	His	Glu	Ala	Tyr	Glu	Ala	65	70	75	80
Tyr	Lys	Ala	Gly	Arg	Ala	Pro	Thr	Pro	Glu	Asp	Phe	Pro	Arg	Gln	Leu	85	90	95	
Ala	Leu	Ile	Lys	Glu	Leu	Val	Asp	Leu	Leu	Gly	Leu	Xaa	Arg	Leu	Glu	100	105	110	
Val	Pro	Gly	Tyr	Glu	Ala	Asp	Asp	Val	Leu	Ala	Thr	Leu	Ala	Lys	Lys	115	120	125	
Ala	Glu	Lys	Glu	Gly	Tyr	Glu	Val	Arg	Ile	Leu	Thr	Ala	Asp	Arg	Asp	130	135	140	
Leu	Tyr	Gln	Leu	Leu	Ser	Asp	Arg	Ile	Ala	Val	Leu	His	Pro	Glu	Gly	145	150	155	160
Tyr	Leu	Ile	Thr	Pro	Ala	Trp	Leu	Trp	Glu	Lys	Tyr	Gly	Leu	Arg	Pro	165	170	175	
Glu	Gln	Trp	Val	Asp	Tyr	Arg	Ala	Leu	Xaa	Gly	Asp	Pro	Ser	Asp	Asn	180	185	190	
Leu	Pro	Gly	Val	Lys	Gly	Ile	Gly	Glu	Lys	Thr	Ala	Xaa	Lys	Leu	Leu	195	200	205	
Xaa	Glu	Trp	Gly	Ser	Leu	Glu	Asn	Leu	Leu	Lys	Asn	Leu	Asp	Arg	Val	210	215	220	
Lys	Pro	Xaa	Xaa	Arg	Glu	Lys	Ile	Xaa	Ala	His	Met	Glu	Asp	Leu	Xaa	225	230	235	240
Leu	Ser	Xaa	Xaa	Leu	Ser	Xaa	Val	Arg	Thr	Asp	Leu	Pro	Leu	Glu	Val	245	250	255	

Asp Phe Ala Xaa Arg Arg Glu Pro Asp Arg Glu Gly Leu Arg Ala Phe  
 260 265 270  
 Leu Glu Arg Leu Glu Phe Gly Ser Leu Leu His Glu Phe Gly Leu Leu  
 275 280 285  
 Glu Xaa Pro Lys Ala Leu Glu Glu Ala Pro Trp Pro Pro Glu Gly  
 290 295 300  
 Ala Phe Val Gly Phe Val Leu Ser Arg Pro Glu Pro Met Trp Ala Glu  
 305 310 315 320  
 Leu Leu Ala Leu Ala Ala Ala Arg Xaa Gly Arg Val His Arg Ala Xaa  
 325 330 335  
 Asp Pro Leu Xaa Gly Leu Arg Asp Leu Lys Glu Val Arg Gly Leu Leu  
 340 345 350  
 Ala Lys Asp Leu Ala Val Leu Ala Leu Arg Glu Gly Leu Asp Leu Xaa  
 355 360 365  
 Pro Gly Asp Asp Pro Met Leu Leu Ala Tyr Leu Leu Asp Pro Ser Asn  
 370 375 380  
 Thr Thr Pro Glu Gly Val Ala Arg Arg Tyr Gly Gly Glu Trp Thr Glu  
 385 390 395 400  
 Asp Ala Gly Glu Arg Ala Leu Leu Ser Glu Arg Leu Phe Xaa Asn Leu  
 405 410 415  
 Xaa Xaa Arg Leu Glu Gly Glu Glu Arg Leu Leu Trp Leu Tyr Xaa Glu  
 420 425 430  
 Val Glu Lys Pro Leu Ser Arg Val Leu Ala His Met Glu Ala Thr Gly  
 435 440 445  
 Val Arg Leu Asp Val Ala Tyr Leu Gln Ala Leu Ser Leu Glu Val Ala  
 450 455 460  
 Glu Glu Ile Arg Arg Leu Glu Glu Glu Val Phe Arg Leu Ala Gly His  
 465 470 475 480  
 Pro Phe Asn Leu Asn Ser Arg Asp Gln Leu Glu Arg Val Leu Phe Asp  
 485 490 495  
 Glu Leu Gly Leu Pro Ala Ile Gly Lys Thr Glu Lys Thr Gly Lys Arg  
 500 505 510  
 Ser Thr Ser Ala Ala Val Leu Glu Ala Leu Arg Glu Ala His Pro Ile  
 515 520 525  
 Val Glu Lys Ile Leu Gln Tyr Arg Glu Leu Thr Lys Leu Lys Asn Thr  
 530 535 540  
 Tyr Ile Asp Pro Leu Pro Xaa Leu Val His Pro Arg Thr Gly Arg Leu  
 545 550 555 560  
 His Thr Arg Phe Asn Gln Thr Ala Thr Ala Thr Gly Arg Leu Ser Ser  
 565 570 575  
 Ser Asp Pro Asn Leu Gln Asn Ile Pro Val Arg Thr Pro Leu Gly Gln  
 580 585 590  
 Arg Ile Arg Arg Ala Phe Val Ala Glu Glu Gly Trp Xaa Leu Val Ala  
 595 600 605

Leu Asp Tyr Ser Gln Ile Glu Leu Arg Val Leu Ala His Leu Ser Gly  
 610 615 620  
 Asp Glu Asn Leu Ile Arg Val Phe Gln Glu Gly Arg Asp Ile His Thr  
 625 630 635 640  
 Gln Thr Ala Ser Trp Met Phe Gly Val Pro Pro Glu Ala Val Asp Pro  
 645 650 655  
 Leu Met Arg Arg Ala Ala Lys Thr Ile Asn Phe Gly Val Leu Tyr Gly  
 660 665 670  
 Met Ser Ala His Arg Leu Ser Gln Glu Leu Ala Ile Pro Tyr Glu Glu  
 675 680 685  
 Ala Val Ala Phe Ile Glu Arg Tyr Phe Gln Ser Phe Pro Lys Val Arg  
 690 695 700  
 Ala Trp Ile Glu Lys Thr Leu Glu Glu Gly Arg Arg Arg Gly Tyr Val  
 705 710 715 720  
 Glu Thr Leu Phe Gly Arg Arg Arg Tyr Val Pro Asp Leu Asn Ala Arg  
 725 730 735  
 Val Lys Ser Val Arg Glu Ala Ala Glu Arg Met Ala Phe Asn Met Pro  
 740 745 750  
 Val Gln Gly Thr Ala Ala Asp Leu Met Lys Leu Ala Met Val Lys Leu  
 755 760 765  
 Phe Pro Arg Leu Xaa Glu Met Gly Ala Arg Met Leu Leu Gln Val His  
 770 775 780  
 Asp Glu Leu Val Leu Glu Ala Pro Lys Xaa Arg Ala Glu Xaa Val Ala  
 785 790 795 800  
 Ala Leu Ala Lys Glu Val Met Glu Gly Val Tyr Pro Leu Ala Val Pro  
 805 810 815  
 Leu Glu Val Glu Val Gly Xaa Gly Glu Asp Trp Leu Ser Ala Lys Glu  
 820 825 830

Xaa

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1647 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATGAATTCGG GGATGCTGCC CCTCTTTGAG CCCAAGGGCC GGGTCCTCCT GGTGGACGGC	60
CACCACCTGG CCTACCGCAC CTTCCACGCC CTGAAGGGCC TCACCACCAG CCGGGGGGAG	120
CCGGTGCAGG CGGTCTACGG CTTGCCAAG AGCCTCCTCA AGGCCCTCAA GGAGGACGGG	180
GACGCGGTGA TCGTGGTCTT TGACGCCAAG GCCCCCTCCT TCCGCCACGA GGCCTACGGG	240
GGGTACAAGG CGGGCCGGGC CCCACGCCG GAGGACTTTC CCCGGCAACT CGCCCTCATC	300
AAGGAGCTGG TGGACCTCCT GGGGCTGGCG CGCCTCGAGG TCCCGGGCTA CGAGGCGGAC	360

GACGTCCTGG CCAGCCTGGC CAAGAAGGCG GAAAAGGAGG GCTACGAGGT CCGCATCCTC	420
ACCGCCGACA AAGACCTTTA CCAGCTCCTT TCCGACCGCA TCCACGTCCT CCACCCCGAG	480
GGGTACCTCA TCACCCCGGC CTGGCTTTGG GAAAAGTACG GCCTGAGGCC CGACCAGTGG	540
GCCGACTACC GGGCCCTGAC CGGGGACGAG TCCGACAACC TTCCCGGGGT CAAGGGCATC	600
GGGGAGAAGA CGGCGAGGAA GCTTCTGGAG GAGTGGGGGA GCCTGGAAGC CCTCCTCAAG	660
AACCTGGACC GGCTGAAGCC CGCCATCCGG GAGAAGATCC TGGCCACAT GGACGATCTG	720
AAGCTCTCCT GGGACCTGGC CAAGGTGCGC ACCGACCTGC CCCTGGAGGT GGACTTCGCC	780
AAAAGGCGGG AGCCCGACCG GGAGAGGCTT AGGGCCTTTC TGGAGAGGCT TGAGTTTGGC	840
AGCCTCCTCC ACGAGTTCGG CCTTCTGGAA AGCCCAAGG CCCTGGAGGA GGCCCCCTGG	900
CCCCCGCCGG AAGGGGCCTT CGTGGGCTTT GTGCTTTCCC GCAAGGAGCC CATGTGGGCC	960
GATCTTCTGG CCCTGGCCGC CGCCAGGGGG GGCCGGGTCC ACCGGGCCCC CGAGCCTTAT	1020
AAAGCCCTCA GGGACCTGAA GGAGGCGCGG GGGCTTCTCG CCAAAGACCT GAGCGTTCTG	1080
GCCCTGAGGG AAGGCCTTGG CCTCCCGCCC GGCACGACC CCATGCTCCT CGCCTACCTC	1140
CTGGACCCTT CCAACACCAC CCCCAGAGGG GTGGCCCGGC GCTACGGCGG GGAGTGGACG	1200
GAGGAGGCGG GGGAGCGGGC CGCCCTTTCC GAGAGGCTCT TCGCCAACCT GTGGGGGAGG	1260
CTTGAGGGGG AGGAGAGGCT CCTTTGGCTT TACCGGGAGG TGGAGAGGCC CCTTTCCGCT	1320
GTCCTGGCCC ACATGGAGGC CACGGGGGTG CGCCTGGACG TGGCCTATCT CAGGGCCTTG	1380
TCCCTGGAGG TGGCCGGGGA GATCGCCCGC CTCGAGGCCG AGGTCTTCCG CCTGGCCGGC	1440
CACCCCTTCA ACCTCAACTC CCGGACCAG CTGGAAAGGG TCCTCTTTGA CGAGCTAGGG	1500
CTTCCCGCCA TCGGCAAGAC GGAGAAGACC GGCAAGCGCT CCACCAGCGC CGCCGTCCTG	1560
GAGGCCCTCC GCGAGGCCCA CCCCATCGTG GAGAAGATCC TGCAGGCATG CAAGCTTGGC	1620
ACTGGCCGTC GTTTTACAAC GTCGTGA	1647

## (2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2088 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ATGAATTCGG GGATGCTGCC CCTCTTTGAG CCAAGGGGCC GGGTCCTCCT GGTGGACGGC	60
CACCACCTGG CCTACCGCAC CTTCACGCC CTGAAGGGCC TCACCACCAG CCGGGGGGAG	120
CCGGTGCAGG CGGTCTACGG CTTCGCCAAG AGCCTCCTCA AGGCCCTCAA GGAGGACGGG	180
GACGCGGTGA TCGTGGTCTT TGACGCCAAG GCCCCCTCCT TCCGCCACGA GGCCTACGGG	240
GGGTACAAGG CGGGCCGGGC CCCACGCCG GAGGACTTTC CCCGGCAACT CGCCCTCATC	300
AAGGAGCTGG TGGACCTCCT GGGGCTGGCG CGCCTCGAGG TCCCGGGCTA CGAGGCGGAC	360

GACGTCCTGG	CCAGCCTGGC	CAAGAAGGCG	GAAAAGGAGG	GCTACGAGGT	CCGCATCCTC	420
ACCGCCGACA	AAGACCTTTA	CCAGCTCCTT	TCCGACCGCA	TCCACGTCCT	CCACCCCGAG	480
GGGTACCTCA	TCACCCCGGC	CTGGCTTTGG	GAAAAGTACG	GCCTGAGGCC	CGACCAGTGG	540
GCCGACTACC	GGGCCCTGAC	CGGGGACGAG	TCCGACAACC	TTCCCGGGGT	CAAGGGCATC	600
GGGGAGAAGA	CGGCGAGGAA	GCTTCTGGAG	GAGTGGGGGA	GCCTGGAAGC	CCTCCTCAAG	660
AACCTGGACC	GGCTGAAGCC	CGCCATCCGG	GAGAAGATCC	TGGCCCACAT	GGACGATCTG	720
AAGCTCTCCT	GGGACCTGGC	CAAGGTGCGC	ACCGACCTGC	CCCTGGAGGT	GGACTTCGCC	780
AAAAGGCGGG	AGCCCGACCG	GGAGAGGCTT	AGGGCCTTTC	TGGAGAGGCT	TGAGTTTGGC	840
AGCCTCCTCC	ACGAGTTCGG	CCTTCTGGAA	AGCCCCAAGG	CCCTGGAGGA	GGCCCCCTGG	900
CCCCCGCCGG	AAGGGGCCTT	CGTGGGCTTT	GTGCTTTCCC	GCAAGGAGCC	CATGTGGGCC	960
GATCTTCTGG	CCCTGGCCGC	CGCCAGGGGG	GGCCGGGTCC	ACCGGGCCCC	CGAGCCTTAT	1020
AAAGCCCTCA	GGGACCTGAA	GGAGGCGCGG	GGGCTTCTCG	CCAAAGACCT	GAGCGTTCTG	1080
GCCCTGAGGG	AAGGCCTTGG	CCTCCCGCCC	GGCGACGACC	CCATGCTCCT	CGCCTACCTC	1140
CTGGACCCTT	CCAACACCAC	CCCCGAGGGG	GTGGCCCGGC	GCTACGGCGG	GGAGTGGACG	1200
GAGGAGGCGG	GGGAGCGGGC	CGCCCTTTCC	GAGAGGCTCT	TCGCCAACCT	GTGGGGGAGG	1260
CTTGAGGGGG	AGGAGAGGCT	CCTTTGGCTT	TACCGGGAGG	TGGAGAGGCC	CCTTTCCGCT	1320
GTCCTGGCCC	ACATGGAGGC	CACGGGGGTG	CGCCTGGACG	TGGCCTATCT	CAGGGCCTTG	1380
TCCCTGGAGG	TGGCCGGGGA	GATCGCCCGC	CTCGAGGCCG	AGGTCTTCCG	CCTGGCCGGC	1440
CACCCCTTCA	ACCTCAACTC	CCGGGACCAG	CTGGAAAGGG	TCCTCTTTGA	CGAGCTAGGG	1500
CTTCCCGCCA	TCGGCAAGAC	GGAGAAGACC	GGCAAGCGCT	CCACCAGCGC	CGCCGTCCTG	1560
GAGGCCCTCC	GCGAGGCCCA	CCCCATCGTG	GAGAAGATCC	TGCAGTACCG	GGAGCTCACC	1620
AAGCTGAAGA	GCACCTACAT	TGACCCCTTG	CCGGACCTCA	TCCACCCCAG	GACGGGCCGC	1680
CTCCACACCC	GCTTCAACCA	GACGGCCACG	GCCACGGGCA	GGCTAAGTAG	CTCCGATCCC	1740
AACCTCCAGA	ACATCCCCGT	CCGCACCCCG	CTTGGGCAGA	GGATCCGCCG	GGCCTTCATC	1800
GCCGAGGAGG	GGTGGCTATT	GGTGGCCCTG	GACTATAGCC	AGATAGAGCT	CAGGGTGCTG	1860
GCCCACCTCT	CCGGCGACGA	GAACCTGATC	CGGGTCTTCC	AGGAGGGGCG	GGACATCCAC	1920
ACGGAGACCG	CCAGCTGGAT	GTTGCGCGTC	CCCCGGGAGG	CCGTGGACCC	CCTGATGCGC	1980
CGGGCGGCCA	AGACCATCAA	CTTCGGGGTC	CTCTACGGCA	TGTCGGCCCA	CCGCCTCTCC	2040
CAGGAGCTAG	CTAGCCATCC	CTTACGAGGA	GGCCCAGGCC	TTCATTGA		2088

## (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 962 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)



## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATGAATTCGG GGATGCTGCC CCTCTTTGAG CCCAAGGGCC GGGTCCTCCT GGTGGACGGC	60
CACCACCTGG CCTACCGCAC CTTCCACGCC CTGAAGGGCC TCACCACCAG CCGGGGGGAG	120
CCGGTGCAGG CGGTCTACGG CTTGCGCAAG AGCCTCCTCA AGGCCCTCAA GGAGGACGGG	180
GACGCGGTGA TCGTGGTCTT TGACGCCAAG GCCCCCTCCT TCCGCCACGA GGCCTACGGG	240
GGGTACAAGG CGGGCCGGGC CCCCACGCCG GAGGACTTTC CCCGGCAACT CGCCCTCATC	300
AAGGAGCTGG TGGACCTCCT GGGGCTGGCG CGCCTCGAGG TCCCGGGCTA CGAGGCGGAC	360
GACGTCCTGG CCAGCCTGGC CAAGAAGGCG GAAAAGGAGG GCTACGAGGT CCGCATCCTC	420
ACCGCCGACA AAGACCTTTA CCAGCTTCTT TCCGACCGCA TCCACGTCCT CCACCCCGAG	480
GGGTACCTCA TCACCCCGGC CTGGCTTTGG GAAAAGTACG GCCTGAGGCC CGACCAGTGG	540
GCCGACTACC GGGCCCTGAC CGGGGACGAG TCCGACAACC TTCCCGGGGT CAAGGGCATC	600
GGGGAGAAGA CGGCGAGGAA GCTTCTGGAG GAGTGGGGGA GCCTGGAAGC CCTCCTCAAG	660
AACCTGGACC GGCTGAAGCC CGCCATCCGG GAGAAGATCC TGGCCACAT GGACGATCTG	720
AAGCTCTCCT GGGACCTGGC CAAGGTGCGC ACCGACCTGC CCCTGGAGGT GGACTTCGCC	780
AAAAGGCGGG AGCCCGACCG GGAGAGGCTT AGGGCCTTTC TGGAGAGGCT TGAGTTTGGC	840
AGCCTCCTCC ACGAGTTCGG CTTCTGGAA AGCCCCAAGT CATGGAGGGG GTGTATCCCC	900
TGGCCGTGCC CCTGGAGGTG GAGGTGGGGA TAGGGGAGGA CTGGCTCTCC GCCAAGGAGT	960
GA	962

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1600 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

ATGGAATTCG GGGATGCTGC CCCTCTTTGA GCCAAGGGC CGGGTCCTCC TGGTGGACGG	60
CCACCACCTG GCCTACCGCA CCTTCCACGC CCTGAAGGGC CTCACCACCA GCCGGGGGGA	120
GCCGGTGCAG GCGGTCTACG GCTTCGCCAA GAGCCTCCTC AAGGCCCTCA AGGAGGACGG	180
GGACGCGGTG ATCGTGGTCT TTGACGCCAA GGCCCCCTCC TTCCGCCACG AGGCCTACGG	240
GGGGTACAAG GCGGGCCGGG CCCCACGCC GGAGGACTTT CCCGGCAAC TCGCCCTCAT	300
CAAGGAGCTG GTGGACCTCC TGGGGCTGGC GCGCCTCGAG GTCCCGGGCT ACGAGGCGGA	360
CGACGTCCTG GCCAGCCTGG CCAAGAAGGC GGAAAAGGAG GGCTACGAGG TCCGCATCCT	420
CACCGCCGAC AAAGACCTTT ACCAGCTCCT TTCCGACCGC ATCCACGTCC TCCACCCCGA	480
GGGGTACCTC ATCACCCCGG CCTGGCTTTG GGAAAAGTAC GGCCTGAGGC CCGACCAGTG	540
GGCCGACTAC CGGGCCCTGA CCGGGGACGA GTCCGACAAC CTTCCCGGGG TCAAGGGCAT	600

CGGGGAGAAG	ACGGCGAGGA	AGCTTCTGGA	GGAGTGGGGG	AGCCTGGAAG	CCCTCCTCAA	660
GAACCTGGAC	CGGCTGAAGC	CCGCCATCCG	GGAGAAGATC	CTGGCCCACA	TGGACGATCT	720
GAAGCTCTCC	TGGGACCTGG	CCAAGGTGCG	CACCGACCTG	CCCCTGGAGG	TGGACTTCGC	780
CAAAAGGCGG	GAGCCCGACC	GGGAGAGGCT	TAGGGCCTTT	CTGGAGAGGC	TTGAGTTTGG	840
CAGCCTCCTC	CACGAGTTTC	GCCTTCTGGA	AAGCCCCAAG	ATCCGCCGGG	CCTTCATCGC	900
CGAGGAGGGG	TGGCTATTGG	TGGCCCTGGA	CTATAGCCAG	ATAGAGCTCA	GGGTGCTGGC	960
CCACCTCTCC	GGCGACGAGA	ACCTGATCCG	GGTCTTCCAG	GAGGGGCGGG	ACATCCACAC	1020
GGAGACCGCC	AGCTGGATGT	TCGGCGTCCC	CCGGGAGGCC	GTGGACCCCC	TGATGCGCCG	1080
GGCGGCCAAG	ACCATCAACT	TCGGGGTCCT	CTACGGCATG	TCGGCCCACC	GCCTCTCCCA	1140
GGAGCTAGCC	ATCCCTTACG	AGGAGGCCCA	GGCCTTCATT	GAGCGCTACT	TTCAGAGCTT	1200
CCCCAAGGTG	CGGGCCTGGA	TTGAGAAGAC	CCTGGAGGAG	GGCAGGAGGC	GGGGGTACGT	1260
GGAGACCCTC	TTCGGCCGCC	GCCGCTACGT	GCCAGACCTA	GAGGCCCGGG	TGAAGAGCGT	1320
GCGGGAGGCG	GCCGAGCGCA	TGGCCTTCAA	CATGCCCGTC	CGGGGCACCG	CCGCCGACCT	1380
CATGAAGCTG	GCTATGGTGA	AGCTCTTCCC	CAGGCTGGAG	GAAATGGGGG	CCAGGATGCT	1440
CCTTCAGGTC	CACGACGAGC	TGGTCCTCGA	GGCCCCAAAA	GAGAGGGCGG	AGGCCGTGGC	1500
CCGGCTGGCC	AAGGAGGTCA	TGGAGGGGGT	GTATCCCCTG	GCCGTGCCCC	TGGAGGTGGA	1560
GGTGGGGATA	GGGGAGGACT	GGCTCTCCGC	CAAGGAGTGA			1600

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CACGAATTCG	GGGATGCTGC	CCCTCTTTGA	GCCCAA	36
------------	------------	------------	--------	----

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 34 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GTGAGATCTA	TCACTCCTTG	GCGGAGAGCC	AGTC	34
------------	------------	------------	------	----

## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 91 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TAATACGACT CACTATAGGG AGACCGGAAT TCGAGCTCGC CCGGGCGAGC TCGAATTCCG 60  
TGTATTCTAT AGTGTACCT AAATCGAATT C 91

## (2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TAATACGACT CACTATAGGG 20

## (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GAATTCGATT TAGGTGACAC TATAGAA 27

## (2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 31 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GTAATCATGG TCATAGCTGG TAGCTTGCTA C 31

## (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 42 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GGATCCTCTA GAGTCGACCT GCAGGCATGC CTACCTTGGT AG

42

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GGATCCTCTA GAGTCGACCT GCAGGCATGC

30

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2502 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

ATGAATTCGG GGATGCTGCC CCTCTTTGAG CCCAAGGGCC GGGTCCTCCT GGTGGACGGC	60
CACCACCTGG CCTACCGCAC CTTCCACGCC CTGAAGGGCC TCACCACCAG CCGGGGGGAG	120
CCGGTGCAGG CGGTCTACGG CTTGCGCAAG AGCCTCCTCA AGGCCCTCAA GGAGGACGGG	180
GACGCGGTGA TCGTGGTCTT TGACGCCAAG GCCCCCTCCT TCCGCCACGA GGCCTACGGG	240
GGGTACAAGG CGGGCCGGGC CCCACGCCG GAGGACTTTC CCCGGCAACT CGCCCTCATC	300
AAGGAGCTGG TGGACCTCCT GGGGCTGGCG CGCCTCGAGG TCCCGGGCTA CGAGGCGGAC	360
GACGTCCTGG CCAGCCTGGC CAAGAAGGCG GAAAAGGAGG GCTACGAGGT CCGCATCCTC	420
ACCGCCGACA AAGACCTTTA CCAGCTCCTT TCCGACCGCA TCCACGTCCT CCACCCCGAG	480
GGGTACCTCA TCACCCCGGC CTGGCTTTGG GAAAAGTACG GCCTGAGGCC CGACCAGTGG	540
GCCGACTACC GGGCCCTGAC CGGGGACGAG TCCGACAACC TTCCCGGGGT CAAGGGCATC	600
GGGGAGAAGA CGGCGAGGAA GCTTCTGGAG GAGTGGGGGA GCCTGGAAGC CCTCCTCAAG	660
AACCTGGACC GGCTGAAGCC CGCCATCCGG GAGAAGATCC TGGCCCACAT GGACGATCTG	720
AAGCTCTCCT GGGACCTGGC CAAGGTGCGC ACCGACCTGC CCCTGGAGGT GGACTTCGCC	780
AAAAGGCGGG AGCCCGACCG GGAGAGGCTT AGGGCCTTTC TGGAGAGGCT TGAGTTTGGC	840
AGCCTCCTCC ACGAGTTCGG CCTTCTGGAA AGCCCCAAGG CCCTGGAGGA GGCCCCCTGG	900
CCCCCGCCGG AAGGGGCCTT CGTGGGCTTT GTGCTTTCCC GCAAGGAGCC CATGTGGGCC	960
GATCTTCTGG CCCTGGCCGC CGCCAGGGGG GGCCGGGTCC ACCGGGCCCC CGAGCCTTAT	1020
AAAGCCCTCA GGGACCTGAA GGAGGCGCGG GGGCTTCTCG CCAAAGACCT GAGCGTTCTG	1080
GCCCTGAGGG AAGGCCTTGG CCTCCCGCCC GGCGACGACC CCATGCTCCT CGCCTACCTC	1140

CTGGACCCTT CCAACACCAC CCCCAGAGGGG GTGGCCCGGC GCTACGGCGG GGAGTGGACG	1200
GAGGAGGCGG GGGAGCGGGC CGCCCTTTCC GAGAGGCTCT TCGCCAACCT GTGGGGGAGG	1260
CTTGAGGGGG AGGAGAGGCT CCTTTGGCTT TACCGGGAGG TGGAGAGGCC CCTTTCCGCT	1320
GTCCTGGCCC ACATGGAGGC CACGGGGGTG CGCCTGGACG TGGCCTATCT CAGGGCCTTG	1380
TCCCTGGAGG TGGCCGGGGA GATCGCCCGC CTCGAGGCCG AGGTCTTCCG CCTGGCCGGC	1440
CACCCCTTCA ACCTCAACTC CCGGGACCAG CTGGAAAGGG TCCTCTTTGA CGAGCTAGGG	1500
CTTCCCGCCA TCGGCAAGAC GGAGAAGACC GGCAAGCGCT CCACCAGCGC CGCCGTCCTG	1560
GAGGCCCTCC GCGAGGCCCA CCCCATCGTG GAGAAGATCC TGCAGTACCG GGAGCTCACC	1620
AAGCTGAAGA GCACCTACAT TGACCCCTTG CCGGACCTCA TCCACCCAG GACGGGCCGC	1680
CTCCACACCC GCTTCAACCA GACGGCCACG GCCACGGGCA GGCTAAGTAG CTCCGATCCC	1740
AACCTCCAGA ACATCCCCGT CCGCACCCCG CTTGGGCAGA GGATCCGCCG GGCCTTCATC	1800
GCCGAGGAGG GGTGGCTATT GGTGGCCCTG GACTATAGCC AGATAGAGCT CAGGGTGCTG	1860
GCCCACCTCT CCGGCGACGA GAACCTGATC CGGGTCTTCC AGGAGGGGCG GGACATCCAC	1920
ACGGAGACCG CCAGCTGGAT GTTCGGCGTC CCCCAGGAGG CCGTGGACCC CCTGATGCGC	1980
CGGGCGGCCA AGACCATCAA CTTCGGGGTC CTCTACGGCA TGTCGGCCCA CCGCCTCTCC	2040
CAGGAGCTAG CCATCCCTTA CGAGGAGGCC CAGGCCTTCA TTGAGCGCTA CTTTCAGAGC	2100
TTCCCCAAGG TCGGGCCTG GATTGAGAAG ACCCTGGAGG AGGGCAGGAG GCGGGGGTAC	2160
GTGGAGACCC TCTTCGGCCG CCGCCGCTAC GTGCCAGACC TAGAGGCCCG GGTGAAGAGC	2220
GTGCGGGAGG CGGCCGAGCG CATGGCCTTC AACATGCCCG TCCGGGGCAC CGCCGCCGAC	2280
CTCATGAAGC TGGCTATGGT GAAGCTCTTC CCCAGGCTGG AGGAAATGGG GGCCAGGATG	2340
CTCCTTCAGG TCCACGACGA GCTGGTCCTC GAGGCCCCAA AAGAGAGGGC GGAGGCCGTG	2400
GCCCGGCTGG CCAAGGAGGT CATGGAGGGG GTGTATCCCC TGGCCGTGCC CCTGGAGGTG	2460
GAGGTGGGGA TAGGGGAGGA CTGGCTCTCC GCCAAGGAGT GA	2502

## (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GATTTAGGTG AACTATAG

19

## (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 72 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CGGACGAACA AGCGAGACAG CGACACAGGT ACCACATGGT ACAAGAGGCA AGAGAGACGA 60  
CACAGCAGAA AC 72

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GTTTCTGCTG TGTCGTCTCT CTTGCCTCTT GTACCATGTG GTACCTGTGT CGCTGTCTCG 60  
CTTGTTTCGTC 70

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GACGAACAAG CGAGACAGCG 20

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GTTTCTGCTG TGTCGTCTCT CTTG 24

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

CCTCTTGTAC CATGTGGTAC CTGTGTCGCT GTCTCGCTTG TTCGTC 46



## (2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 50 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

ACACAGGTAC CACATGGTAC AAGAGGCAAG AGAGACGACA CAGCAGAAAC

50

## (2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 15 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Met Ala Ser Met Thr Gly Gly Gln Gln Met Gly Arg Ile Asn Ser  
 1 5 10 15

## (2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 969 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

ATGGCTAGCA TGA CTGGTGG ACAGCAAATG GGTCGGATCA ATTCGGGGAT GCTGCCCCCTC 60  
 TTTGAGCCCA AGGGCCGGGT CCTCCTGGTG GACGGCCACC ACCTGGCCTA CCGCACCTTC 120  
 CACGCCCTGA AGGGCCTCAC CACCAGCCGG GGGGAGCCGG TGCAGGCGGT CTACGGCTTC 180  
 GCCAAGAGCC TCCTCAAGGC CCTCAAGGAG GACGGGGACG CGGTGATCGT GGTCTTTGAC 240  
 GCCAAGGCCC CCTCCTTCCG CCACGAGGCC TACGGGGGGT ACAAGGCGGG CCGGGCCCCC 300  
 ACGCCGGAGG ACTTTCCCCG GCAACTCGCC CTCATCAAGG AGCTGGTGGA CCTCCTGGGG 360  
 CTGGCGCGCC TCGAGGTCCC GGGCTACGAG GCGGACGACG TCCTGGCCAG CCTGGCCAAG 420  
 AAGGCGGAAA AGGAGGGCTA CGAGGTCCGC ATCCTCACCG CCGACAAAGA CCTTTACCAG 480  
 CTTCTTTCCG ACCGCATCCA CGTCCTCCAC CCCGAGGGGT ACCTCATCAC CCCGGCCTGG 540  
 CTTTGGGAAA AGTACGGCCT GAGGCCCGAC CAGTGGGCCG ACTACCGGGC CCTGACCGGG 600  
 GACGAGTCCG ACAACCTTCC CGGGGTCAAG GGCATCGGGG AGAAGACGGC GAGGAAGCTT 660  
 CTGGAGGAGT GGGGGAGCCT GGAAGCCCTC CTCAAGAACC TGGACCGGCT GAAGCCCGCC 720  
 ATCCGGGAGA AGATCCTGGC CCACATGGAC GATCTGAAGC TCTCCTGGGA CCTGGCCAAG 780

GTGCGCACCG ACCTGCCCCCT GGAGGTGGAC TTCGCCAAAA GGCGGGAGCC CGACCGGGAG	840
AGGCTTAGGG CCTTTCTGGA GAGGCTTGAG TTTGGCAGCC TCCTCCACGA GTTCGGCCTT	900
CTGGAAAGCC CCAAGTCATG GAGGGGGTGT ATCCCCTGGC CGTGCCCCTG GAGGTGGAGG	960
TGGGGATAG	969

## (2) INFORMATION FOR SEQ ID NO:31:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 948 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ATGGCTAGCA TGACTGGTGG ACAGCAAATG GGTCGGATCA ATTCGGGGAT GCTGCCCCTC	60
TTTGAGCCCA AGGGCCGGGT CCTCCTGGTG GACGGCCACC ACCTGGCCTA CCGCACCTTC	120
CACGCCCTGA AGGGCCTCAC CACCAGCCGG GGGGAGCCGG TGCAGGCGGT CTACGGCTTC	180
GCCAAGAGCC TCCTCAAGGC CCTCAAGGAG GACGGGGACG CGGTGATCGT GGTCTTTGAC	240
GCCAAGGCCC CCTCCTTCCG CCACGAGGCC TACGGGGGGT ACAAGGCGGG CCGGGCCCCC	300
ACGCCGGAGG ACTTTCCCCG GCAACTCGCC CTCATCAAGG AGCTGGTGGG CCTCCTGGGG	360
CTGGCGCGCC TCGAGGTCCC GGGCTACGAG GCGGACGACG TCCTGGCCAG CCTGGCCAAG	420
AAGGCGGAAA AGGAGGGCTA CGAGGTCCGC ATCCTCACCG CCGACAAAGA CCTTTACCAG	480
CTTCTTTCCG ACCGCATCCA CGTCCTCCAC CCCGAGGGGT ACCTCATCAC CCCGGCCTGG	540
CTTTGGGAAA AGTACGGCCT GAGGCCCGAC CAGTGGGCCG ACTACCGGGC CCTGACCGGG	600
GACGAGTCCG ACAACCTTCC CGGGGTCAAG GGCATCGGGG AGAAGACGGC GAGGAAGCTT	660
CTGGAGGAGT GGGGGAGCCT GGAAGCCCTC CTCAAGAACC TGGACCGGCT GAAGCCCGCC	720
ATCCGGGAGA AGATCCTGGC CCACATGGAC GATCTGAAGC TCTCCTGGGA CCTGGCCAAG	780
GTGCGCACCG ACCTGCCCCCT GGAGGTGGAC TTCGCCAAAA GGCGGGAGCC CGACCGGGAG	840
AGGCTTAGGG CCTTTCTGGA GAGGCTTGAG TTTGGCAGCC TCCTCCAAGA GTTCGGCCTT	900
CTGGAAAGCC CCAAGGCCGC ACTCGAGCAC CACCACCACC ACCACTGA	948

## (2) INFORMATION FOR SEQ ID NO:32:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 206 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

CGCCAGGGTT TTCCAGTCA CGACGTTGTA AAACGACGGC CAGTGAATTG TAATACGACT	60
CACTATAGGG CGAATTCGAG CTCGGTACCC GGGGATCCTC TAGAGTCGAC CTGCAGGCAT	120

GCAAGCTTGA GTATTCTATA GTGTCACCTA AATAGCTTGG CGTAATCATG GTCATAGCTG 180  
TTTCCTGTGT GAAATTGTTA TCCGCT 206

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 43 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

TTCTGGGTTC TCTGCTCTCT GGTCGCTGTC TCGCTTGTTTC GTC 43

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GCTGTCTCGC TTGTTTCGTC 19

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

GACGAACAAG CGAGACAGCG 20

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

TTCTGGGTTC TCTGCTCTCT GGTC 24

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 43 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:  
GACGAACAAG CGAGACAGCG ACCAGAGAGC AGAGAACCCA GAA 43

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 23 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:  
ACCAGAGAGC AGAGAACCCA GAA 23

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:  
AACAGCTATG ACCATGATTA C 21

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 30 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:  
GGATCCTCTA GAGTCGACCT GCAGGCATGC 30

## CLAIMS

We claim:

1. A DNA sequence encoding a thermostable DNA polymerase altered in sequence relative to the native sequence such that it exhibits altered DNA synthetic activity from that of the native DNA polymerase.
2. The DNA sequence of Claim 1 wherein the encoded DNA polymerase exhibits reduced synthetic activity from that of the native DNA polymerase.
3. The DNA sequence of Claim 1 wherein the alteration to said native sequence comprises a change in a single nucleotide.
4. The DNA sequence of Claim 1 wherein the alteration to said native sequence comprises a deletion.
5. The DNA sequence of Claim 4 comprising a DNA sequence selected from the group consisting of SEQ ID NOS:9-12 and 21.
6. The DNA sequence of Claim 1 wherein the alteration to said native sequence comprises an insertion.
7. The DNA sequence of Claim 2 derived from an organism selected from the group consisting of *Thermus aquaticus*, *Thermus flavus* and *Thermus thermophilus*.
8. A recombinant DNA vector comprising a DNA sequence encoding a thermostable DNA polymerase altered in sequence relative to the native sequence such that it exhibits altered DNA synthetic activity from that of the native DNA polymerase.

9. The recombinant DNA vector comprising a DNA sequence of Claim 8 wherein the alteration to said native sequence comprises a change in a single nucleotide.
10. The recombinant DNA vector comprising a DNA sequence of Claim 8 wherein the alteration to said native sequence comprises a deletion.
11. The recombinant DNA vector comprising a DNA sequence of Claim 8 derived from an organism selected from the group consisting of *Thermus aquaticus*, *Thermus flavus* and *Thermus thermophilus*.
12. The recombinant DNA vector comprising a DNA sequence of Claim 11 comprising a DNA sequence selected from the group consisting of SEQ ID NOS:9-12 and 21.
13. A host cell transformed with the recombinant vector of Claim 8.
14. A thermostable DNA polymerase altered in amino acid sequence such that it exhibits altered DNA synthetic activity from that of the native DNA polymerase but retains substantially the same 5' nuclease activity of that of the native DNA polymerase.
15. The polymerase of Claim 14 wherein the altered polymerase exhibits reduced synthetic activity from that of the native DNA polymerase.
16. The polymerase of Claim 15 wherein the alteration to said native sequence comprises a change in an amino acid.
17. The polymerase of Claim 15 wherein the alteration to said native sequence comprises a deletion.



18. The polymerase of Claim 15 derived from an organism selected from the group consisting of *Thermus aquaticus*, *Thermus flavus* and *Thermus thermophilus*.

19. The polymerase of Claim 18 comprising an amino acid sequence encoded by the nucleic acid sequences selected from the group consisting of SEQ ID NOS:9-12 and 21.

20. A method of detecting the presence of a specific target DNA molecule comprising:

a) providing:

i) a target nucleic acid,

ii) a first oligonucleotide complementary to a first portion of said target nucleic acid, and

iii) a second oligonucleotide, a region of which is complementary to a second portion of said target nucleic acid, said non-complementary region of said second oligonucleotide providing a single-stranded arm at its 5' end;

b) mixing said target nucleic acid, said first oligonucleotide and said second oligonucleotide under conditions wherein said first oligonucleotide and the 3' end of said second oligonucleotide are annealed to said target DNA sequence so as to create a first cleavage structure;

c) providing a cleavage means under conditions such that cleavage of said first cleavage structure occurs preferentially at a site located within said second oligonucleotide in a manner dependent upon the annealing of said first and second oligonucleotides on said target nucleic acid, thereby liberating the single-stranded arm of said second oligonucleotide generating a third oligonucleotide;

d) providing a first hairpin structure having a single-stranded 3' arm and a single-stranded 5' arm under conditions wherein said third oligonucleotide anneals to said single-stranded 3' arm of said first hairpin thereby creating a second cleavage structure;

- 5 e) providing conditions under which cleavage of said second cleavage structure occurs by said cleavage means liberating the single-stranded 5' arm of said second cleavage structure so as to create reaction products comprising a fourth oligonucleotide and a first cleaved hairpin detection molecule;
- f) providing a second hairpin structure having a single-stranded 3' arm and a single-stranded 5' arm under conditions wherein said fourth oligonucleotide anneals to the single-stranded 3' arm of said second hairpin thereby creating a third cleavage structure;
- 10 g) providing conditions under which cleavage of said third cleavage structure occurs by said cleavage means liberating the single-stranded 5' arm of said third cleavage structure so as to create reaction products comprising a fifth oligonucleotide identical in sequence to said third oligonucleotide and a second cleaved hairpin detection molecule; and
- 15 h) detecting the presence of said first and second cleaved hairpin detection molecules.

21. The method of Claim 20 wherein steps d) through g) are repeated at least once.

20 22. The method of Claim 20 wherein said cleavage means comprises an altered thermostable DNA polymerase such that cleavage reactions occur in the absence of any significant polymerase activity.

23. The method of Claim 20 wherein the cleavage reactions of steps c), e) and g) will not occur absent the annealing of said first oligonucleotide, said third oligonucleotide and said fourth oligonucleotide, respectively.

25 24. The method of Claim 20 wherein the cleavage reaction of step c) occurs within the annealed portion of said second oligonucleotide.

25. The method of Claim 20 wherein the cleavage reaction of step c) occurs within the non-annealed portion of said second oligonucleotide.

26. A method of detecting the presence of a specific target nucleic acid molecule comprising:

- 5 a) providing:
- i) a cleavage means,
  - ii) a target nucleic acid,
  - iii) a first oligonucleotide complementary to a first portion of said target nucleic acid,
  - 10 iv) a first solid support having a second oligonucleotide, a region of which is complementary to a second portion of said target nucleic acid, said non-complementary region of said second oligonucleotide providing a single-stranded arm at its 5' end, a portion of said 5' arm comprising a first signal oligonucleotide,
  - 15 v) a plurality of uncleaved second solid supports each having a third oligonucleotide, a region of which is complementary to said first signal oligonucleotide, the non-complementary region of said third oligonucleotide providing a single-stranded arm at its 5' end, a portion of said 5' arm comprising a second signal
  - 20 oligonucleotide, and
  - vi) a plurality of uncleaved third solid supports each having a fourth oligonucleotide, a region of which is complementary to said second signal oligonucleotide, the non-complementary region of said fourth oligonucleotide providing a single-stranded arm at its
  - 25 5' end, a portion of said 5' arm comprising said first signal oligonucleotide;
- b) mixing said cleavage means, said target nucleic acid, said first oligonucleotide and said second oligonucleotide under conditions wherein said first oligonucleotide and the 3' end of said second
- 30 oligonucleotide are annealed to said target DNA sequence so as to create a

first cleavage structure and cleavage of said first cleavage structure results in the liberating of said first signal oligonucleotide;

5 c) reacting said liberated first signal oligonucleotide with one of said plurality of second solid supports under conditions such that said first signal oligonucleotide hybridizes to said complementary region of said third oligonucleotide to create a second cleavage structure and cleavage of said second cleavage structure results in the liberating of said second signal oligonucleotide and a cleaved second solid support;

10 d) reacting said liberated second signal oligonucleotide with one of said plurality of third solid supports under conditions such that said second signal oligonucleotide hybridizes to said complementary region of said fourth oligonucleotide to create a third cleavage structure and cleavage of said third cleavage structure results in the liberating of a second molecule of said first signal oligonucleotide and a cleaved third solid support; and


15 e) detecting the presence of said first and second signal oligonucleotides.



27. The method of Claim 26 wherein said cleavage means comprises a 5' nuclease derived from a thermostable DNA polymerase.

20 28. The method of Claim 27 wherein said thermostable DNA polymerase is derived from an organism selected from the group consisting of *Thermus aquaticus*, *Thermus flavus* and *Thermus thermophilus*.

29. The method of Claim 28 wherein said 5' nuclease is encoded by a DNA sequence selected from the group consisting of SEQ ID NOS:11, 30 and 31.

FIGURE 1A



**PART ONE: TRIGGER REACTION****PART TWO: DETECTION REACTION****FIG. 1B**

2/43



FIG. 2A

MAJORITY	ATGXXGGCGATGCITCCCTCITTGAGCCCAAGCGCGGTCTCTGGTGGACGGCCACCACTGGCCT	
DNAPTAQ	...AG..G.....G.....G.....	70
DNAPTFL	.....G.....G.....G.....	67
DNAPTTH	...GA.....G.....A.....	70
MAJORITY	ACCGCACCTTCITCGCCCTGAAGGGCTCACCACGAGCGCGCGGCGAACCCTGCAGCGGCTCTACGGCTT	
DNAPTAQ	.....GA.....G.....G.....	140
DNAPTFL	.....T.....G.....C.....C..T.....	137
DNAPTTH	.....G.....G.....	140
MAJORITY	CGCCAAAGAGCCTCCTCAAGGCCCTGAAGGAGGACGGGACXXGGCGGTGXTGGTGTCTTGACGGCAAG	
DNAPTAQ	.....G.....G.....A.....	207
DNAPTFL	.....A.....GT..T.....	204
DNAPTTH	.....T..AA..C..CT.....	210
MAJORITY	GCCCCCTCCTTCGGCCACGAGGCCCTACGAGGCCCTACAAGCGCGCGCGCGCGCGGAGACTTTC	
DNAPTAQ	.....G..GG.....G.....	277
DNAPTFL	.....G.....G.....	274
DNAPTTH	.....GA.....G.....G.....	280
MAJORITY	CGCGGCAGCTCGCCCTCATCAAGGAGCTGCTGGACCTCCTGGCGCTTGGCGGCTCGAGGTCGCCCGGCTA	
DNAPTAQ	.....A.....G.....G.....	347
DNAPTFL	.....G.....T.....A..C.....T..G..G.....T.....	344
DNAPTTH	.....T.....T..A.C.....	350

FIG. 2B

MAJORITY	CGAGCGGAGCAGGTGCTGGCCACCCCTGGCCCAAGAGGGGAAAGGAGGGGTACGAGGTGGCGCATCCTC	
DNAPTAQ	.....C.....G.....C.....C.....	417
DNAPTFL	T.....G.....CG.....	414
DNAPTTH	.....T..C.....	420
MAJORITY	ACCGCGGACCGGACCTCTACCAAGCTCCCTTCGGACCGCATCGCGGTCCCTCCACCGCGAGGGGTACCTCA	
DNAPTAQ	.....AAA.....T.....CA.....	487
DNAPTFL	..T.....G.....G.....A.....T.....G.	484
DNAPTTH	.....A..G.C.....G.....CC.....	490
MAJORITY	TCACCCCGCGGTGGCTTTGGGAGAAAGTACGGCCTGAGCGCGGAGCAGTGGGTGGACTACCGCGGCCCTGGC	
DNAPTAQ	.....C.....A.....C..C.....CC.....A.	557
DNAPTFL	.....AC.....C.C.....	554
DNAPTTH	.....A.....C.....T..C.....C.T	560
MAJORITY	GGGGACCCCTCCGACAAACCTCCCGGGGTCAAGGGCATCGGGGAGAGACCGCCXGAAGCTCCTCXAG	
DNAPTAQ	C.....GAG.....T.....G..GAG.....T..GG..	627
DNAPTFL	.....G..T..A.....G.....A..G...A..CGG	624
DNAPTTH	.....TC.....A..	630
MAJORITY	GAGTGGCGGAGCCCTGGAAACCTCCTCAAGAACCTGGACCGGTGAAGCCCGC...CXTCGGGAGAGA	
DNAPTAQ	.....GC.....C.....A.....	694
DNAPTFL	.....T..C..C.....A.....T..T..G.....C	691
DNAPTTH	.....A.....A....A.AAAA.G.....	700

FIG. 2C

MAJORITY	TCGAGGGCCGACATGGAXGACCTGAXGCTCTCCTGGAGGCTXTCCGAGGTGGGCACCGACCTGCCCCCTGGA	
DNAPTAO	...T.....C..T....A.....C..GG..A.....	764
DNAPTFL	...GGG.....G.C....GGC..T...C..A....T.....A...T.....	761
DNAPTTH	..A.....C....A.....C.G.....T.....C.....C.....	770
MAJORITY	GGTGGACTTCGCCCAAGXGGGGAGCCGGAGCGGGGAGGGGCTTCTGGAGAGGCTGGAGCTT	
DNAPTAO	.....AA.....A.....A.....T.....T.....	834
DNAPTFL	.....GG.G.C.C..CACA...A...T.....T..GC...T...T.....C..T.....	831
DNAPTTH	.....C.....C..G.....C.....C.....C.....C.....	840
MAJORITY	GGCAGCCTCCTCCACGAGTTCGGCCTCCTGGAGGGCCCCCAAGGCCCTGGAGGAGGCCCCCTGGCCCCCGGC	
DNAPTAO	.....T.....T.....AA.....	904
DNAPTFL	..A.....G.....G.....G..G....GGCA.....T..	901
DNAPTTH	.....C.....C.....GCCC.....	910
MAJORITY	CGGAAGGGGCCCTTCGTGGGCTTTGTCTTTCCGGCCCGAGCCCATGTGGGCCGAGCTTCTGCCCTGGC	
DNAPTAO	.....G.....AAG.....T.....	974
DNAPTFL	.....T..TT.....TC.T.....T.....	971
DNAPTTH	.....C.....C.....G.....AAA.....	980
MAJORITY	CGCCGCCAGGGAGGGCCGGTCCACGGGGCAGCAGACCCCTTAXGGGGCTXAGGGACCTXAAGGAGGTG	
DNAPTAO	.....G.....C..C..G..T.A..AA.C...C.....G.....C..	1044
DNAPTFL	T.GG..GT.....G..CC...T.....A.....C.....G.....T.....G....	1041
DNAPTTH	....TG.....C.....G.....G.....GGC...G..A.A.....C.....C..	1050

**SUBSTITUTE SHEET (RULE 26)**

6/43

FIG. 2E

MAJORITY	GGAGATCCGGCCCTCGAGGAGGAGGTCTTCGGGCTGGCCGGCCACCCGTTCAAGCTCAACTCCCGGGGAC	
DNAPTAQ	.....GC.....CC.....	1464
DNAPTFL	...G.G...AG..G.....	1461
DNAPTTH	.....T.....G.....	1470
MAJORITY	CAGCTGGAAGGGTGGTCTTTGACGAGCTXGGGCTTCGGGCCATCGGCAAGACGGAGAACXGGCAAGC	
DNAPTAQ	.....C.....A.....	1534
DNAPTFL	.....GC.....G..G..T.....	1531
DNAPTTH	.....TA.....T.G..G.....C.A.....	1540
MAJORITY	GCTCCACCGAGCGCGCGGCTGCTGGAGGGCCCTXCGXGAGGGCCGACCCCATCGTGGAGAAAGATCCTGCAGTA	
DNAPTAQ	.....C.....C..C.....	1604
DNAPTFL	.....T.....G..A.....CCGC.....	1601
DNAPTTH	.....G.....A..G.....C...C..	1610
MAJORITY	CCGGGAGCTCACCAAGCTCAAGAACACCTACATXGACCCGCTGCCXGXCCCTCGTCCACCCGAGGACGGGGC	
DNAPTAQ	.....G.....G.....T.....G.A...A.....	1674
DNAPTFL	.....A.....A.....C.C...G...A...C...A...C...	1671
DNAPTTH	.....G.G.....C..AAG.....G.....	1680
MAJORITY	CGGCTCCACACCGGCTTCAACCAGACGGGGCAGGGCCAGGGCTTAGTACCTCCGACCCCAACCTGC	
DNAPTAQ	.....A.....T.....C..	1744
DNAPTFL	..G.....C.....TCC.....	1741
DNAPTTH	.....G.....	1750

FIG. 2F

MAJORITY	AGAACATCCCGCTCCGCACCCXCTGGGCCAGAGGATCCGCCGGGCGTTCGTGGCCGAGGAGCGXTCGGT	
DNAPTAQ	.....G..T..G.....A.C.....G...C.	1814
DNAPTFL	.....G.....T.....C.C.....A.....C.....	1811
DNAPTTH	.....CT.....C.....C...T...C	1820
MAJORITY	GTGGTGGCCCTGGACTATAGCCAGATAGAGCTCCGGGTCCCTGGCCGACCTCTCCGGGGACGAGAACCTG	
DNAPTAQ	A.....T.T.....C.....A...G.....C.....	1884
DNAPTFL	.....T.T.....C.....T.....T.....	1881
DNAPTTH	.....T.....C.....C.....A.....	1890
MAJORITY	ATCCGGGTCITCCAGGAGGGAGGACATCCACAGCCAGACGGCCAGCTGGATGTTCCGGCGTCCCCCGGG	
DNAPTAQ	.....G.....C.....GG.....G..	1954
DNAPTFL	.....T.....A.....A.....TT...C.	1951
DNAPTTH	.....A.....A.....A.....	1960
MAJORITY	AGGCCGTGGACCCCTGATGGCGGGGGCCAGACCATCAACTTCGGGGTCCCTCTACGGCATGTCCGC	
DNAPTAQ	.....A.GG..A...T.....G..	2024
DNAPTFL	.....GG.....GG.G.....G.....	2021
DNAPTTH	.....GG.G.....G.....	2030
MAJORITY	GCACGGCGCTCTCCGAGGAGCTTGGCCATCCCCCTACGAGGCGGGTGGGCTTCATTGAGCGGCTACTTCGAG	
DNAPTAQ	.....A.....T.....CCA.....T...	2094
DNAPTFL	.....GG.....T.....	2091
DNAPTTH	...TA.G.....T..A.....A	2100



FIG. 2G

MAJORITY	AGCTTCCCCCAAGGTGCGGGCCCTGGATTGAGAAAGACCCCTGGAGGGCAGGGCGGTACGTGGAGA	
DNAPTAO	.....	2164
DNAPTFL	...A.....GG.....C.....C.CC.....T.....	2161
DNAPTTH	.....A..A.....G...A.....C.....A.	2170
MAJORITY	CCCTCTTCGGCGCGGGGCTAGCTGCCCGGACCTCAACGGCCGGGTGAAGAGCGTGGCGGAGCGCGGGGA	
DNAPTAO	.....C.....A.....AG.G.....C..	2234
DNAPTFL	.....T.....C.....	2231
DNAPTTH	.....AA.AA.....CA.....C.....	2240
MAJORITY	GCGCATGGGCTTCAACATGCCCGTCCAGGGCACCGCCGGACCTCATGAAGCTGGCCATGGTGAAGCTC	
DNAPTAO	.....T.....	2304
DNAPTFL	.....G.....CG...T	2301
DNAPTTH	.....C.....C.....	2310
MAJORITY	TTCCCGCGGCTXCAGGAAATGGGGCCAGGATGCTCGTXCAGGTCCACGAGCGCTGGTCCCTCGAGGCCCC	
DNAPTAO	.....A...GG.....T.....	2374
DNAPTFL	.....T.....C.....G.....TT.G.....G.....	2371
DNAPTTH	.....C..C.G...G.....C.C.....C.....CC.....G.....	2380
MAJORITY	CCAAAGAGCGGGGAGGXGGTGGCCCGCTTGGCCCAAGGAGGTGATGGAGGGGTCATCCCGCTGGCCGT	
DNAPTAO	.....A.....A.....GG.....GGG.....G.....	2444
DNAPTFL	.....G..C.....AG...A.....GG.....CAG..	2441
DNAPTTH	..C...C.....C...A.....G.....C.....AA..C.....C.....	2450

FIG. 2H

MAJORITY	GGCCCTGGAGGTGGAGGTGGGATGGGGAGGACTGGCTCTCCGCCAAGGAGTAG	2499
DNAPTAA	.....A..... GA	2496
DNAPTFL	.....CC.....	2505
DNAPTTH	.....T..... GT...	

FIG. 3A

MAJORITY	MXA ML PL FEPKGRVLLVDGHHLAYRTFFALKGLTTSRGEPVQAVYGFAKSLKALKEDG·DAVXVVVFD AK	
TAQ PRO	·RG·····H·····I·····	69
TFL PRO	·····V·····V·····	68
TTH PRO	·E·····YK··F·····	70
MAJORITY	AP SFRHEAYEAYKAGRAPTEDFPROLALI KELVDLLGLXRLEVPYCGYEADDVLATLAKKAEKEGYEVRI L	
TAQ PRO	·····GG·····A·····S·····	139
TFL PRO	·····V·····F·····R·····	138
TTH PRO	·····FT·····	140
MAJORITY	TADRDLYQLLSORIAVLHPGEGYLITPAWLWEKYGLRPEQWVDYRALXGDPSONLPGVKGIGECTAXKLLX	
TAQ PRO	·····K·····H·····D··A·····T··E·····R··E	209
TFL PRO	·····E·····I·····Y·····A·····I·····QR·IR	208
TTH PRO	·····V·····V·····H·····E·····F·····V·····L··K	210
MAJORITY	EWG SLENLLKNLDRVKP·XXREKI XAHMEDLXL SXXLSXVRTDLPLEVDFAXRREPDRGLRAFLE RLEF	
TAQ PRO	·····A·····L·····AI·····L·····D··K··WD·AK·····K·····R·····	278
TFL PRO	·····FQH··O·····SL·····LO·G··A··A··RK··O·H·····GR··T·NL·····	277
TTH PRO	·····ENV·····K··L··R··LE··R·····L·QG·····	280
MAJORITY	GSL LHEFGLLEXPKALEEAPWPPPEGAFVGFVLSRPEPMWAE L L A A A R X G R V H R A X D P L X G L R D L K E V	
TAQ PRO	·····S·····D·····K·····D·····G·····PE·YKA·····A	348
TFL PRO	·····G··A·····L··SF·····G·WE··L··O··R·····G·	347
TTH PRO	·····A·AP·····K·····C·D·····A··A··K·····	350

FIG. 3B

MAJORITY	RGLLAKDLAVLALREGLDLXPGDDPMLLAYLLDPSNTTPEGVARRYGGWTE	DAGERALLSERLFXNLXX	
TAQ PRO	.....S.....G.P.....	.....E.....A.....	418
TRL PRO	.....I.....F.E.....	.....A.....OT.KE	417
TTH PRO	.....S.....V.....	.....AH.....HR..LK	420
MAJORITY	RLEGEERLLWLYXEVEKPLSRVLAHMEATGVRLDVAYLOALSLEVAEEI	RRLEEEVFRLAGHPFNLNSRD	
TAQ PRO	.....R...R...A.....	.....R.....A.....	488
TRL PRO	.....K.....E.....R.....	.....EA.V.O.....	487
TTH PRO	.....K.....H.....	.....L.....	490
MAJORITY	QLERVLFDELGLPAIGKTEKTGKRSTSAAVLEALREAHPIVEKI	LQYRELTCLKNTYIDPLPXLVHPRTG	
TAQ PRO	.....	.....S.....D.I.....	558
TRL PRO	.....R...L...O.....	.....DR.....A....K..	557
TTH PRO	.....	.....H.....V.....S.....	560
MAJORITY	RLHTRFNOTATATGRLSSSDPNLQNI	PVRTPLGORIRRAFVAEEGWXLVALDYSOIELRVLAHLSGDENL	
TAQ PRO	.....	.....I.....L.....	628
TRL PRO	.....	.....V...V.....	627
TTH PRO	.....	.....A...A.....	630
MAJORITY	IRVFQEGRDIHTOTASWMFGVPPPEAVDPLMRRAAKTINFGVLYGMSAHL	SOELAIPEEEAVAFIERYFO	
TAQ PRO	.....E.....R.....	.....O.....	698
TRL PRO	.....S...G.....	.....G...S.....	697
TTH PRO	.....K.....V.....	.....	700

FIG. 3C

MAJORITY	SFPKVRAWIEKTL EEGRRRGYVETLFGRRRYVPDLNARVKSUREAAERMAFNMPVOGTAADLMK LAMVKL	
TAQ PRO	.....E.....	768
TFL PRO	Y.....G.....R.	767
TTH PRO	.....K.....	770
MAJORITY	FPRLXEMGARM LQVHDELVL EAPKXRAEXVAALAKEVMEGVYPLAVPLEVEVGXGEDWLSAKEX	
TAQ PRO	.....E.....A..R.....I.....	833
TFL PRO	.....O.L.....D.....R.....W..O.....L.....	831
TTH PRO	.....R.....L.....OA.....E.....A..KA.....M.....G	835

FIGURE 4








FIGURE 5








FIG. 7




FIG. 8

	1	2	3	4	5	6
DNAP-T:	-	+	+	+	+	+
MgCl <sub>2</sub> :	+	-	+	+	+	+
dNTPs:	+	-	+	-	+	-
Primers:	+	-	+	+	-	-




FIG. 9A



FIG. 9B




FIG. 10B

FIG. 10A






FIG. 11A

FIG. 11B




FIG. 12A




FIG. 13A




FIG. 12B




FIG. 13B

FIGURE 14






FIGURE 15



P<sub>φ10</sub>: Bacteriophage T7 φ10 promoter  
Tφ: T7 φ Terminator  
RBS: Ribosome binding site

FIGURE 16








FIG. 17




FIG. 18





FIG. 19A

	"4d"	NO POL. ACTIVITY	"4b" 2 PT. MUTATION SMALL ACTIVITY	UNMODIFIED DNAP Tag	dNTP
1	2	3	4	5	6
				7	8
				-	+
				-	+







FIG. 20A


SEQUENCE OF ALPHA PRIMER:

5' GAC GAA CAA GCG AGA CAG CG 3'

**FIG. 20B**



CLEAVED A - HAIRPIN



CLEAVED T - HAIRPIN

**FIG. 20C**

TOP = T - HAIRPIN  
 BOTTOM = A - HAIRPIN

Rsa I  
 Mnl I  
 Nla III  
 HgiC I  
 Nla IV  
 Rsa I  
 Kpn I  
 BsmAI

GTTTCGTGTCGTCCTCTCTTGCCCTCTCTTGACCATGTGGTACCTGTGCTGCTGCTGCTTGTTCGTC  
 CAAAGACGACACAGCAGAGAACGGAGAACATGGTACACCATGGACACAGCGACAGAGCGAACACAGCAGGC

72

**FIG. 20D**



FIGURE 21





FIGURE 22A







FIG. 22B

FIGURE 23



CDR BEAD	T	T	T	AT	AT	A	A	A
PILOT	-	-	+	+	-	+	+	-
CLEAVASE	M	M	+	+	+	+	+	-
	M						M	M




FIG. 24




FIG. 25B

FIG. 25A









FIGURE 27



**FIG. 28A**

**FIG. 28B**

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/06253

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) : Please See Extra Sheet.

US CL : 435/6, 91.1, 172.3, 252.1, 320.1; 530/350; 536/23.7

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.1, 172.3, 252.1, 320.1; 530/350; 536/23.7

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, BIOSIS, CA

search terms: synthesis, polymerization, polymerase, deficient, thermostable, Taq, lack

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP, A, 482714 (SULLIVAN) 29 April 1992, Abstract.	1, 3
Y	Nucleic Acids Research, Volume 18, No. 24, issued 1990, M. J. Longley et al, "Characterization of the 5' to 3' exonuclease associated with <u>Thermus aquaticus</u> DNA polymerase", pages 7317-7322, especially page 7320, Substrate Specificity and page 7319, second column.	1-19

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance		
"E" earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed	"&"	document member of the same patent family

Date of the actual completion of the international search

25 AUGUST 1994

Date of mailing of the international search report

SEP 06 1994

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DAVID SCHREIBER

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/06253

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Journal of Biological Chemistry, Volume 264, No. 11, issued 15 April 1989, F. C. Lawyer et al, "Isolation, Characterization, and Expression in <u>Escherichia coli</u> of the DNA Polymerase Gene from <u>Thermus aquaticus</u> ", pages 6427-6437, especially Figures 2 and 7, Materials and Methods, and page 6434, second full paragraph.	1-19
Y	Proceedings of the National Academy of Science USA, Volume 88, issued August 1991, P. M. Holland et al, "Detection of specific polymerase chain reaction product by utilizing the 5'-->3' exonuclease activity of <u>Thermus aquaticus</u> DNA polymerase", pages 7276-7280, especially page 7276, paragraph bridging first and second column and Figure 1.	1-19
Y	Proceedings of the National Academy of Science USA, Volume 82, issued January 1985, T. A. Kunkel, "Rapid and efficient site-specific mutagenesis without phenotypic selection", pages 488-492, especially Abstract.	1-19

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/06253

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Telephone Practice

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/06253

## A. CLASSIFICATION OF SUBJECT MATTER: IPC (5):

C12Q 1/68; C12P 19/34; C12N 1/21, 9/16, 15/63, 15/70; C07H 21/02, 21/04

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

- I. Claims 1-13, drawn to DNA encoding a thermostable polymerase altered in nucleotide sequence such that it exhibits altered DNA synthetic activity from that of the native enzyme, a vector carrying the DNA, and a host carrying the recombinant vector, classified in Class 536, subclass 23.7, for example.
- II. Claims 14-25, drawn to a thermostable polymerase altered in nucleotide sequence such that it exhibits altered DNA synthetic activity from that of the native enzyme and a method of detecting the presence of a specific target DNA molecule utilizing complementary and partially complementary oligonucleotides which form hairpin cleavage structures and the altered thermostable polymerase, classified in Class 530, subclass 350, for example.
- III. Claims 26-29, drawn to a method of detecting the presence of a specific target DNA molecule utilizing support bound partially complementary oligonucleotides which form cleavage structures which are not hairpin structures, classified in Class 435, subclass 6.